For Research Use Only

uPAR/CD87 Polyclonal antibody

Catalog Number: 10286-1-AP 26 Publications



Basic Information

Catalog Number: 10286-1-AP Concentration: 400 ug/ml

UNIPROT ID: Source: Rabbit Q03405 Full Name: Isotype: **PLAUR**

Immunogen Catalog Number: Calculated MW: AG0187 37 kDa

Observed MW: 35-70 kDa

BC002788

5329

GeneID (NCBI):

GenBank Accession Number:

Purification Method:

Antigen affinity purification

Recommended Dilutions: WB: 1:1000-1:8000 IF/ICC: 1:200-1:800

Applications

Tested Applications: WB, IF/ICC, ELISA Cited Applications: WB, IHC, IF, CoIP Species Specificity: human, mouse

Cited Species: human, mouse, bovine Positive Controls:

WB: RAW 264.7 cells, A2780 cells, U-251 cells, U-937

IF/ICC: U-251 cells,

Background Information

uPAR is a 45-65 kDa, highly glycosylated, GPI-anchored membrane protein. In addition to the membrane-anchored form, uPAR is released from the plasma membrane by cleavage of the GPI anchor and can be found as a soluble form (suPAR). uPAR contains three homologous domains (D1-D3) of which the N-terminal one (D1) represents the uPAbinding domain. After binding to uPAR, uPA cleaves plasminogen, generating the active protease plasmin which is involved in a wide variety of physiologic and pathologic processes. In addition to regulating proteolysis, uPAR has important function in cell adhesion, migration and proliferation. Studies reveal that uPAR expression is elevated during inflammation and tissue remodelling and in many human cancers, in which it frequently indicates poor prognosis. (PMID 20027185; 12461559)

Notable Publications

| Author | Pubmed ID | Journal | Application |
|----------------|-----------|-----------------|-------------|
| Mulin Liu | 34552345 | J Inflamm Res | IHC |
| Jun-Ming Huang | 30245631 | Front Pharmacol | IF |
| Min Liu | 28871079 | Nat Commun | WB |

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 10286-1-AP (uPAR/CD87 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Ethanol) fixed U-251 cells using uPAR/CD87 antibody (10286-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2).