For Research Use Only

Lactotransferrin/LTF Polyclonal antibody

Catalog Number: 10933-1-AP

Featured Product

13 Publications



Basic Information

Catalog Number: 10933-1-AP Concentration: 600 ug/ml

Source:
Rabbit
Isotype:
IgG
Immunogen Catalog Number:

AG1379

Observed MW: 30,51,80 kDa

GenBank Accession Number:

BC015822 GeneID (NCBI): 4057

UNIPROT ID: P02788 Full Name: lactotransferrin

Calculated MW: 80 kDa

Positive Controls:

WB: human saliva, HEK-293 cells, human pancreas

Purification Method:

WB: 1:1000-1:4000

protein lysate

IHC: 1:20-1:200

Antigen affinity purification

IP: 0.5-4.0 ug for 1.0-3.0 mg of total

Recommended Dilutions:

tissue

IP : HeLa cells,

IHC: human liver tissue, human prostate hyperplasia

tissue

Applications

Tested Applications: WB, IHC, IP, ELISA Cited Applications: WB, IHC, IF Species Specificity: human

Cited Species: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

LTF(Lactotransferrin) is also named as GIG12, LF and belongs to the transferrin family. It is an iron binding transport protein which can bind two atoms of ferric iron in association with the binding of an anion, usually bicarbonate. LTF is also an important protein component of the innate immune system that is broadly distributed within the body fluids. It can contribute to the activation of both the innate and adaptive immune responses by promoting the recruitment of leukocytes and activation of dendritic cells(PMID:18453607). This protein has 2 isoforms produced by alternative promoter usage. The human LTF can be isolated two fragments, an N-terminal tryptic fragment having an MW of 30 kDa and a C-terminal tryptic fragment having an MW of 50 kDa by a mild tryptic digest(PMID:3790094).

Notable Publications

Author	Pubmed ID	Journal	Application
Qiong Huang	36185743	Bioact Mater	WB
Rujia Qin	34266418	Cancer Cell Int	WB,IHC
Kaoru Araki-Sasaki	30214321	Int Med Case Rep J	IHC

Storage

Storage

Store at -20°C. Stable for one year after shipment.

Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 4006900926 E: Proteintech-CN@ptglab.com

W: ptgcn.co

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




human saliva were subjected to SDS PAGE followed by western blot with 10933-1-AP (Lactoferrin antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human liver using 10933-1-AP (Lactoferrin antibody) at dilution of 1:50 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human liver using 10933-1-AP (Lactoferrin antibody) at dilution of 1:50 (under 40x lens).



IP result of anti-LTF (IP:10933-1-AP, 4ug; Detection:10933-1-AP 1:600) with HeLa cells lysate 4000ug.