### For Research Use Only

# HNRNPC Polyclonal antibody

Catalog Number: 11760-1-AP

Featured Product

45 Publications



**Basic Information** 

Catalog Number: GenBank Accession Number: 11760-1-AP BC003394 GeneID (NCBI): Concentration: 500 μg/ml 3183 **UNIPROT ID:** Source: Rabbit P07910 Full Name: Isotype:

heterogeneous nuclear ribonucleoprotein C (C1/C2) Immunogen Catalog Number:

AG2356 Calculated MW:

> 32 kDa Observed MW: 32 kDa, 36-40 kDa

**Purification Method:** 

Antigen affinity purification Recommended Dilutions:

WB: 1:5000-1:50000

IP: 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC: 1:250-1:1000 IF/ICC: 1:300-1:1200

**Applications** 

**Tested Applications:** WB, IHC, IF/ICC, IP, ELISA

Cited Applications: WB, IHC, IF, IP, RIP Species Specificity: human, mouse **Cited Species:** 

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

human, mouse

Positive Controls:

WB: HeLa cells, human brain tissue, K-562 cells, HepG2

cells, MCF-7 cells IP: HeLa cells.

IHC: mouse colon tissue, human lymphoma tissue

IF/ICC: HeLa cells,

# **Background Information**

 $HNRNPC, also \ named \ as \ Heterogeneous \ nuclear \ ribonucleoproteins \ C1/C2, is \ a \ 306 \ amino \ acid \ protein, which \ and \ an \ acid \ protein, which \ acid \ protein \ acid \ acid$ contains 1 RRM (RNA recognition motif) domain and belongs to the RRM HNRPC family. HNRNPC localizes in the nucleus. It binds pre-mRNA and nucleates the assembly of 40S hnRNP particles. HNRNPC may play a role in the early steps of spliceosome assembly and pre-mRNA splicing. HNRNPC can act as a tetramer and is involved in the assembly of 405 hnRNP particles and plays a role with CSDE1 in internal initiation of translation during mitosis.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Li Wang	36255739	Hum Mol Genet	WB
Li-Chong Wang	33134160	Front Oncol	WB,IHC
Fanhua Kong	36434140	Discov Oncol	WB

Storage

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3


Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926 E: Proteintech-CN@ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## **Selected Validation Data**




WB result of HNRNPC antibody (11760-1-AP; 1:20000; incubated at room temperature for 1.5 hours) with sh-Control and sh-HNRNPC transfected HeLa cells.




Various lysates were subjected to SDS PAGE followed by western blot with 11760-1-AP (HNRNPC antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



IP result of anti-HNRNPC (IP:11760-1-AP, 4ug; Detection:11760-1-AP 1:10000) with HeLa cells lysate 1360 ug.



Immunohistochemical analysis of paraffinembedded mouse colon tissue slide using 11760-1-AP (HNRNPC antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using HNRNPC antibody (11760-1-AP) at dilution of 1:600 and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red).