For Research Use Only

Caspase 4/P20/P10 Polyclonal antibody

www.ptglab.com

Purification Method:

WB: 1:500-1:1000 IHC: 1:20-1:200

IF/ICC: 1:200-1:800

Antigen affinity purification

Recommended Dilutions:

Catalog Number: 11856-1-AP

Featured Product

16 Publications

Basic Information

Catalog Number: GenBank Accession Number: 11856-1-AP BC017839 Concentration: GeneID (NCBI): 600 ug/ml **UNIPROT ID:** Source: Rabbit P49662 Full Name: Isotype:

caspase 4, apoptosis-related cysteine

Immunogen Catalog Number: AG2429 Calculated MW: 377 aa. 43 kDa

Observed MW: 43-48 kDa

peptidase

Applications

Tested Applications: WB, IHC, IF/ICC, ELISA **Cited Applications:** WB, IHC, IF Species Specificity:

human **Cited Species:** human, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

IF/ICC: HepG2 cells,

WB: SGC-7901 cells, THP-1 cells IHC: human pancreas cancer tissue,

Background Information

Caspase 4 (CASP4), is a member of the cysteine-aspartic acid protease (caspase) family. Sequential activation of caspases plays a central role in the execution-phase of cell apoptosis. Caspase 4 is able to cleave and activate its own precursor protein, as well as caspase 1 precursor. Overexpression of caspase 4 will induce cell apoptosis.

Notable Publications

Author	Pubmed ID	Journal	Application
Liangmin Fu	36248876	Front Immunol	IHC,WB
Xiao-Lei Xin	31542790	Med Sci Monit	WB
Si Wen	34494385	J Diabetes Investig	


Storage

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




SGC-7901 cells were subjected to SDS PAGE followed by western blot with 11856-1-AP (Caspase 4/P20/P10 antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human pancreas cancer using 11856-1-AP (Caspase 4/P20/P10 antibody) at dilution of 1:50 (under 25x lens).



Various lysates were subjected to SDS PAGE followed by western blot with 11856-1-AP (Caspase 4/P20/P10 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



SGC-7901 cells were subjected to SDS PAGE followed by western blot with 11856-1-AP (Caspase 4/P20/P10 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using Caspase 4/P20/P10 antibody (11856-1-AP) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2).