

# Glutamate receptor 2 Polyclonal antibody

Catalog Number: 11994-1-AP

49 Publications

## Basic Information

## Catalog Number:

11994-1-AP

## Concentration:

600 ug/ml

## Source:

Rabbit

## Isotype:

IgG

## Immunogen Catalog Number:

AG2662

## GenBank Accession Number:

BC010574

## GeneID (NCBI):

2891

## UNIPROT ID:

P42262

## Full Name:

glutamate receptor, ionotropic, AMPA 2

## Calculated MW:

883 aa, 99 kDa

## Observed MW:

94-105 kDa

## Purification Method:

Antigen affinity purification

## Recommended Dilutions:

WB: 1:2000-1:12000

IP: 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

IHC: 1:50-1:500

IF-P: 1:50-1:500

## Applications

## Tested Applications:

WB, IHC, IF-P, IP, ELISA

## Cited Applications:

WB, IHC, IF, IP, CoIP

## Species Specificity:

human, mouse, rat

## Cited Species:

human, mouse, rat

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

## Positive Controls:

WB : mouse brain tissue, rat brain tissue

IP : mouse brain tissue,

IHC : mouse brain tissue,

IF-P : mouse eye tissue,

## Background Information

Glutamate receptors are the predominant excitatory neurotransmitter receptors in the mammalian brain and are activated in a variety of normal neurophysiologic processes. GRIA2 (glutamate receptor 2, GluR-2) belongs to a family of glutamate receptors that are sensitive to alpha-amino-3-hydroxy-5-methyl-4-isoxazole propionate (AMPA), and function as ligand-activated cation channels that mediate the fast component of excitatory postsynaptic currents in neurons of the central nervous system. These channels are assembled from 4 related subunits, GRIA1-4. The presence of GRIA2 in the heteromeric AMPA receptors impermeabilizes it to calcium, preventing possible calcium-mediated toxicity.

## Notable Publications

Author	Pubmed ID	Journal	Application
Pengcheng Ma	36179027	Sci Adv	WB, CoIP
YuanYuan Hou	34710402	Exp Neurol	WB
Joanna Gruszczynska-Biegala	27826230	Front Cell Neurosci	WB

## Storage

## Storage:

Store at -20°C. Stable for one year after shipment.

## Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 4006900926


E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)W: [ptgcn.com](http://ptgcn.com)

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 11994-1-AP (Glutamate receptor 2 antibody) at dilution of 1:6000 incubated at room temperature for 1.5 hours.




IP result of anti-Glutamate receptor 2 (IP:11994-1-AP, 4ug; Detection:11994-1-AP 1:2000) with mouse brain tissue lysate 2100 ug.



Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 11994-1-AP (Glutamate receptor 2 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed mouse eye tissue using Glutamate receptor 2 antibody (11994-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded mouse eye tissue using Glutamate receptor 2 antibody (11994-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).