

For Research Use Only

# MFN2 Polyclonal antibody

Catalog Number: 12186-1-AP

Featured Product

449 Publications



## Basic Information

### Catalog Number:

12186-1-AP

### Concentration:

500 ug/ml

### Source:

Rabbit

### Isotype:

IgG

### Immunogen Catalog Number:

AG2845

### GenBank Accession Number:

BC017061

### GeneID (NCBI):

9927

### ENSEMBL Gene ID:

ENSG00000116688

### UNIPROT ID:

O95140

### Full Name:

mitofusin 2

### Calculated MW:

757 aa, 86 kDa

### Observed MW:

86 kDa

### Purification Method:

Antigen affinity purification

### Recommended Dilutions:

WB: 1:5000-1:50000

IP: 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

IHC: 1:50-1:500

## Applications

### Tested Applications:

WB, IHC, IP, ELISA

### Cited Applications:

WB, IHC, IF, IP, CoIP

### Species Specificity:

human, mouse, rat

### Cited Species:

human, mouse, rat, pig, monkey, chicken, zebrafish, bovine, goat, duck

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

### Positive Controls:

**WB** : mouse brain tissue, HeLa cells, mouse kidney tissue, rat brain tissue, rat heart tissue, mouse liver tissue, rat kidney tissue, rat liver tissue

**IP** : mouse kidney tissue,

**IHC** : human colon cancer tissue, mouse kidney tissue

## Background Information

MFN2, also named as CPRP1 and KIAA0214, belongs to the mitofusin family. It is an Essential transmembrane GTPase, which mediates mitochondrial fusion. MFN2 acts independently of the cytoskeleton. It therefore plays a central role in mitochondrial metabolism and may be associated with obesity and/or apoptosis processes. Overexpression of MFN2 induces the formation of mitochondrial networks. It plays an important role in the regulation of vascular smooth muscle cell proliferation. Defects in MFN2 are the cause of Charcot-Marie-Tooth disease type 2A2 (CMT2A2). Defects in MFN2 are the cause of Charcot-Marie-Tooth disease type 6 (CMT6). Ubiquitinated forms of Mfn2 (mono- and polyubiquitinated) are present during mitophagy.

## Notable Publications

Author	Pubmed ID	Journal	Application
Maria Manczak	27677309	Hum Mol Genet	IF
Na Jiang	32975326	Cell Prolif	WB,IHC
Siwen Li	28957766	Chemosphere	WB

## Storage

### Storage:

Store at -20°C. Stable for one year after shipment.

### Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)


W: [ptgcn.com](http://ptgcn.com)

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 12186-1-AP (MFN2 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.




Various lysates were subjected to SDS PAGE followed by western blot with 12186-1-AP (MFN2 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.




WB result of MFN2 antibody (12186-1-AP; 1:1000; incubated at room temperature for 1.5 hours) with sh-Control and sh-MFN2 transfected HeLa cells.




Immunohistochemical analysis of paraffin-embedded human colon cancer tissue slide using 12186-1-AP (MFN2 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP result of anti-MFN2 (IP:12186-1-AP, 4ug; Detection:12186-1-AP 1:10000) with mouse kidney tissue lysate 2160 ug.



Immunohistochemical analysis of paraffin-embedded human colon cancer tissue slide using 12186-1-AP (MFN2 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse kidney tissue slide using 12186-1-AP (MFN2 antibody) at dilution of 1:800 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).