For Research Use Only

ETFA Polyclonal antibody

Catalog Number: 12262-1-AP 7 Publications



Basic Information

Catalog Number: GenBank Accession Number: 12262-1-AP BC015526 GeneID (NCBI): Concentration: 450 µg/ml 2108 **UNIPROT ID:** Source: Rabbit P13804 Full Name: Isotype:

100 µl suspension electron-transfer-flavoprotein, alpha polypeptide

AG2931 Calculated MW: 30 kDa Observed MW: 30 kDa

Applications

Tested Applications:

WB, IHC, IF/ICC, FC (Intra), ELISA

Immunogen Catalog Number:

Cited Applications:

Species Specificity: human, mouse, rat **Cited Species:** human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: Y79 cells, human ileum tissue, mouse small intestine tissue, rat small intestine tissue

Purification Method:

WB: 1:1000-1:3000 IHC: 1:50-1:500

IF/ICC: 1:50-1:500

Antigen affinity purification

FC (Intra): 0.80 ug per 10^6 cells in a

Recommended Dilutions:

IHC: human breast cancer tissue,

IF/ICC : HeLa cells, FC (Intra): MCF-7 cells,

Background Information

Electron transfer flavoprotein (ETF) exists in the mitochondrial matrix as a heterodimer of 30 kD alpha subunits (ETFA) and 28 kD beta subunits (ETFB) and contains 1 flavin adenine dinucleotide (FAD) and 1 adenosine 5-prime monophosphate (AMP) per heterodimer. ETFA serves as a specific electron acceptor for several dehydrogenases, including five acyl-CoA dehydrogenases, glutaryl-CoA and sarcosine dehydrogenase.

Notable Publications

Author	Pubmed ID	Journal	Application
Bin Jiang	35381197	Mol Cell	IF
Min-Zhi Peng	35727412	Cell Mol Life Sci	WB
Singaravelu Ragunath R	20181094	Proteome Sci	WB

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 12262-1-AP (ETFA antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 12262-1-AP (ETFA antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using ETFA antibody (12262-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L), 594phalloidin (red).



1x10^6 MCF-7 cells were intracellularly stained with 0.8 ug ETFA Polyclonal antibody (12262-1-AP) and Coralite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2)(red), or 0.8 ug Rabbit IgG control Rabbit PolyAb (30000-0-AP) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).