

For Research Use Only

# Alix Polyclonal antibody

Catalog Number: 12422-1-AP

Featured Product

264 Publications



## Basic Information

### Catalog Number:

12422-1-AP

### Concentration:

800 ug/ml

### Source:

Rabbit

### Isotype:

IgG

### Immunogen Catalog Number:

AG3074

### GenBank Accession Number:

BC020066

### GeneID (NCBI):

10015

### UNIPROT ID:

Q8WUM4

### Full Name:

programmed cell death 6 interacting protein

### Calculated MW:

868 aa, 96 kDa

### Observed MW:

75-100 kDa

### Purification Method:

Antigen affinity purification

### Recommended Dilutions:

WB: 1:5000-1:50000

IP: 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

IHC: 1:50-1:500

IF/ICC: 1:200-1:800

## Applications

### Tested Applications:

WB, IHC, IF/ICC, IP, ELISA

### Cited Applications:

WB, IHC, IF, IP

### Species Specificity:

human, mouse, rat

### Cited Species:

human, mouse, rat, pig, rabbit, monkey, hamster

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

### Positive Controls:

**WB**: HEK-293 cells, human urine exosomes tissue, NIH/3T3 cells, HeLa cells, Jurkat cells, mouse liver tissue, rat liver tissue

**IP**: Jurkat cells,

**IHC**: human liver cancer tissue, human colon tissue

**IF/ICC**: HeLa cells,

## Background Information

ALG-2-interacting protein 1 (ALIX), also known as AIP1 or Hp95, is encoded by PDCD6IP gene and is involved in cell death through mechanisms involving its binding partner ALG-2 (apoptosis-linked gene-2). ALG-2 is a 22-kDa protein containing five serially repetitive EF-hand structures and is defined as a regulator of calcium-induced apoptosis following endoplasmic reticulum (ER) stress. ALIX interacts with ALG-2 through its C-terminal proline-rich region and participates in formation of multivesicular bodies. Recent finding suggest that ALIX is a critical component of caspase 9 activation and apoptosis triggered by calcium. The alix antibody recognizes an additional band of 75-80 kDa which has also been observed in cells and exosomes.

## Notable Publications

Author	Pubmed ID	Journal	Application
Christopher R Silvers	34564696	Br J Cancer	WB
Denghui Wei	32958903	Cell Res	WB
Xue Zou	33002418	Mol Ther	WB

## Storage

### Storage:

Store at -20°C. Stable for one year after shipment.

### Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)


W: [ptgcn.com](http://ptgcn.com)

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 12422-1-AP (Alix antibody) at dilution of 1:15000 incubated at room temperature for 1.5 hours.




human urine exosome samples were subjected to SDS PAGE followed by western blot with 12422-1-AP (Alix antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.




IP result of anti-Alix (IP:12422-1-AP, 3ug; Detection:12422-1-AP 1:500) with Jurkat cells lysate 4000ug.



WB result of Alix antibody (12422-1-AP; 1:8000; incubated at room temperature for 1.5 hours) with sh-Control and sh-Alix transfected HEK-293 cells.



Immunohistochemical analysis of paraffin-embedded human liver cancer tissue slide using 12422-1-AP (Alix antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Methanol) fixed HeLa cells using Alix antibody (12422-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).