## For Research Use Only

## GLRA2 Polyclonal antibody

Catalog Number: 13831-1-AP



**Basic Information** 

Catalog Number: GenBank Accession Number: 13831-1-AP BC032864 GeneID (NCBI): Size: 300 μg/ml 2742 **UNIPROT ID:** Source:

Rabbit P23416 Full Name: Isotype:

glycine receptor, alpha 2

Calculated MW: Immunogen Catalog Number: 452 aa, 52 kDa Observed MW:

48 kDa

**Purification Method:** Antigen affinity purification Recommended Dilutions: WB 1:500-1:1000 IHC 1:20-1:200

**Applications** 

**Tested Applications:** IHC, WB, ELISA Species Specificity:

human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

WB: PBMC cells, K-562 cells IHC: mouse brain tissue,

## **Background Information**

Glycine is an inhibitory neurotransmitter in the central nervous system. The glycine receptor (GlyR) is a member of the nicotinic acetylcholine receptor of ligand-gated ion channels. This receptor has important roles in a variety of physiological processes, especially in mediating inhibitory neurotransmission in the spinal cord and brain stem. GlyR is a pentameric receptor comprised of alpha and beta subunits. Four alpha-subunits (GLRA1, GLRA2, GLRA3, GLRA4) and one beta-subunit (GLRB) of GlyR have been identified. (PMID: 11409700; 15383648)

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**




WB result of GLRA2 antibody (13831-1-AP, 1:600) with PBMC cells. Proteintech  $\alpha$  GLRA2 13831-1-AP in 10% BSA / TBS 0.05% Tween-20 0/N @ 4°C Goat anti-rabbit HRP 1:10000 in 5% milk / TBS 0.05% Tween-20 1h @ RT.



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 13831-1-AP (GLRA2 Antibody) at dilution of 1:50 (under 10x lens).



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 13831-1-AP (GLRA2 Antibody) at dilution of 1:50 (under 40x lens).



Various lysates were subjected to SDS PAGE followed by western blot with 13831-1-AP (GLRA2 antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.