## For Research Use Only

## ZAP70 Polyclonal antibody

Catalog Number: 15592-1-AP

Featured Product

1 Publications

BC039039

GenBank Accession Number:



**Basic Information** 

Catalog Number: 15592-1-AP

GeneID (NCBI): Concentration: 750 ug/ml

ENSEMBL Gene ID: Source: Rabbit ENSG00000115085 **UNIPROT ID:** Isotype: P43403

Immunogen Catalog Number: Full Name:

AG7943 zeta-chain (TCR) associated protein

kinase 70kDa Calculated MW: 70 kDa Observed MW: 70 kDa

**Purification Method:** 

Antigen affinity purification Recommended Dilutions:

WB: 1:5000-1:20000

IP: 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

IHC: 1:20-1:200

**Applications** 

**Tested Applications:** WB, IHC, IP, ELISA Cited Applications:

Species Specificity: human, mouse, rat Cited Species:

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: Jurkat cells, A375 cells

IP: Jurkat cells,

IHC: human tonsillitis tissue, human kidney tissue, human lymphoma tissue, human testis tissue

**Background Information** 

Zeta-associated protein (ZAP)-70, a member of the Syk family of tyrosine kinases, plays an important role in T-cell receptor signaling, natural killer (NK) cell activation and early B-cell development. ZAP-70 is expressed in a subset of cases of chronic lymphocytic leukemia/small lymphocytic lymphoma (CLL/SLL) with unmutated immunoglobulin heavy-chain variable region (IgVH) genes and is associated with poor clinical outcome.

**Notable Publications** 

Author Pubmed ID Journal Application J Immunol Methods WB Jiaqi Zheng 38484791

Storage


Storage:

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

## **Selected Validation Data**




WB result of ZAP70 antibody (15592-1-AP, 1:20000) with si-control and si-ZAP70 transfected Jurkat cells.




Jurkat cells were subjected to SDS PAGE followed by western blot with 15592-1-AP (ZAP70 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human tonsillitis using 15592-1-AP (ZAP70 antibody) at dilution of 1:50 (under 10x lens)



Immunohistochemical analysis of paraffinembedded human tonsillitis using 15592-1-AP (ZAP70 antibody) at dilution of 1:50 (under 40x lens).



IP result of anti-ZAP70 (IP:15592-1-AP, 4ug; Detection:15592-1-AP 1:1000) with Jurkat cells lysate 3200ug.



Immunofluorescent analysis of (4% PFA) fixed RAW 264.7 cells using ZAP70 antibody (15592-1-AP) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).