For Research Use Only

MFF Polyclonal antibody

Catalog Number: 17090-1-AP

Featured Product

201 Publications



Basic Information

Catalog Number:

GenBank Accession Number:

17090-1-AP

BC000797

Concentration:

350 ug/ml

56947

Source:

UNIPROT ID:

Rabbit

Q9GZY8

Isotype:

GenBank Accession Number:

BC000797

UNCBI):

350 ug/ml

56947

UNIPROT ID:

Q9GZY8

Full Name:

gG mitochondrial fission factor

Immunogen Catalog Number:Calculated MW:AG999038 kDa

Observed MW: 35-38 kDa Purification Method: Antigen affinity purification

Recommended Dilutions:

WB: 1:5000-1:50000 IP: 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC: 1:500-1:2000 IF/ICC: 1:50-1:500

Applications

Tested Applications: WB, IHC, IF/ICC, IP, ELISA

Cited Applications: WB, IHC, IF, IP Species Specificity: human, mouse, rat Cited Species:

human, mouse, rat, pig, monkey, hamster, goat, squirrel

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: mouse brain tissue, rat brain tissue, mouse heart

tissue, rat heart tissue IP: mouse brain tissue,

IHC: human stomach tissue, mouse brain tissue

IF/ICC: HepG2 cells,

Background Information

MFF (mitochondrial fission factor) is a mitochondrial outer membrane protein that is involved in mitochondrial localization of Drp1 and mitochondrial fission. Multiple isoforms of MFF exist due to the alternative splicing. This antibody recognizes the endogenous MFF protein around 26-29 kDa and 35-38 kDa.

Notable Publications

Author	Pubmed ID	Journal	Application
Katherine Labbé	34586346	J Cell Biol	WB
Q E Xie	34571250	Mitochondrion	WB
Feixiang Bao	30237514	Cell Death Dis	WB

Storage

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Immunohistochemical analysis of paraffinembedded human stomach tissue slide using 17090-1-AP (MFF antibody) at dilution of 1:1000 (under 10x lens).



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using MFF antibody (17090-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Various lysates were subjected to SDS PAGE followed by western blot with 17090-1-AP (MFF antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.



IP result of anti-MFF (IP:17090-1-AP, 4ug; Detection:17090-1-AP 1:30000) with mouse brain tissue lysate 1680 ug.