For Research Use Only

RIPK1-Specific Polyclonal antibody

Catalog Number: 17519-1-AP

Featured Product

136 Publications



Basic Information

Catalog Number:

17519-1-AP

Concentration:

800 ug/ml

Source:

Rabbit

Listype:

GenBank Accession Number:

NM_003804

GeneID (NCBI):

8737

UNIPROT ID:

Q13546

Full Name:

receptor (TNFRSF)-interacting serine-

Calculated MW: 76 kDa Observed MW: 70-80 kDa

threonine kinase 1

Applications

Tested Applications:
WB, IHC, ELISA
Cited Applications:
WB, IHC, IF, IP, CoIP
Species Specificity:
mouse, rat
Cited Species:

human, mouse, rat, pig, yeast

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: mouse spleen tissue, rat liver tissue, rat heart

Purification Method:

WB: 1:1000-1:8000 IHC: 1:50-1:500

Antigen affinity purification

Recommended Dilutions:

tissue, rat spleen tissue

IHC: mouse small intestine tissue.

Background Information

RIPK1(Receptor-interacting serine/threonine-protein kinase 1) is primarily involved in mediating TNF-R1-induced cell activation, apoptosis, and necroptosis and belongs to a novel class of kinases that function in cell survival and cell death mechanisms(PMID:22685397). It has 2 isoforms produced by alternative splicing. It also can exist as a protein with a lower molecular weight of about 25 kDa and 45 kDa in HAEC and HUVEC(PMID:22685397).

Notable Publications

Author	Pubmed ID	Journal	Application
Xingkang Wu	34597653	J Ethnopharmacol	WB
Ying Tu	36155221	Environ Pollut	WB
Xiao-Ling Zhou	33041798	Front Pharmacol	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3 $\,$

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 4006900926 E: Proteintech-CN@ptglab.com


W: ptgcn.co

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 17519-1-AP (RIPK1-Specific antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded mouse small intestine tissue slide using 17519-1-AP (RIPK1-Specific antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).