# For Research Use Only

# CD133 Polyclonal antibody

Catalog Number: 18470-1-AP

Featured Product

262 Publications

BC012089

8842

043490 Full Name:

97 kDa

GeneID (NCBI):

**UNIPROT ID:** 

prominin 1 Calculated MW:

GenBank Accession Number:



**Basic Information** 

Catalog Number: 18470-1-AP Concentration:

1100 µg/ml Source: Rabbit Isotype: Immunogen Catalog Number:

AG13328

Observed MW: 115 kDa-120 kDa **Purification Method:** 

Antigen affinity purification Recommended Dilutions: WB: 1:2000-1:10000

IHC: 1:4000-1:16000 IF-P: 1:50-1:500

**Applications** 

**Tested Applications:** WB, IHC, IF-P, ELISA Cited Applications: WB, IHC, IF Species Specificity:

human, mouse, rat **Cited Species:** human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

#### **Positive Controls:**

WB: HT-29 cells, mouse kidney tissue, mouse pancreas tissue, rat kidney tissue, Caco-2 cells

IHC: mouse kidney tissue, IF-P: mouse kidney tissue,

# **Background Information**

CD133, also known as PROM1 (prominin-1) or AC133, belongs to the prominin family. CD133 is a transmembrane glycoprotein with an NH2-terminal extracellular domain, five transmembrane loops and a cytoplasmic tail. The expression of CD133 has been reported in hematopoietic stem cells, endothelial progenitor cells, neuronal and glial stem cells, suggesting the potential role of CD133 as a cell surface marker of adult stem cells. CD133 has also been reported as a marker of cancer stem cells in various human tumors.

### **Notable Publications**

Author	Pubmed ID	Journal	Application
Yang Liu	27685621	Cell Death Dis	WB,IF
Qiu-Zhong Pan	25267273	Stem Cells	WB, IHC
Chenlong Li	31558707	Cell Death Dis	WB

# Storage


Storage:

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


# **Selected Validation Data**




Various lysates were subjected to SDS PAGE followed by western blot with 18470-1-AP (CD133 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded mouse kidney tissue slide using 18470-1-AP (CD133 antibody) at dilution of 1:8000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse kidney tissue slide using 18470-1-AP (CD133 antibody) at dilution of 1:8000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed mouse kidney tissue using CD133 antibody (18470-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L).