

For Research Use Only

CD117/c-Kit Polyclonal antibody

Catalog Number: 18696-1-AP

35 Publications



Basic Information

Catalog Number:

18696-1-AP

Concentration:

550 ug/ml

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

BC071593

GeneID (NCBI):

3815

UNIPROT ID:

P10721

Full Name:

v-kit Hardy-Zuckerman 4 feline
sarcoma viral oncogene homolog

Calculated MW:

110 kDa

Observed MW:

140-160 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB: 1:1000-1:4000

IHC: 1:50-1:500

IF/ICC: 1:50-1:500

Applications

Tested Applications:

WB, IHC, IF/ICC, ELISA

Cited Applications:

WB, IHC, IF

Species Specificity:

human

Cited Species:

human, mouse, rat, sheep, goat

**Note-IHC: suggested antigen retrieval with
TE buffer pH 9.0; (*) Alternatively, antigen
retrieval may be performed with citrate
buffer pH 6.0**

Positive Controls:

WB : Mo7e cells,

IHC : stromal tumor tissue, human placenta tissue,
human testis tissue, Insulinoma tissue

IF/ICC : MCF-7 cells,

Background Information

CD117, also known as c-Kit and SCFR, is a transmembrane protein with tyrosine kinase activity encoded by the oncogene c-kit (PMID: 2448137). It is a member of the type III receptor tyrosine kinase family, which also includes CSF-1R, PDGFR β , PDGFR α , and FLT3 (PMID: 29518044). CD117 is expressed on hematopoietic stem cells and progenitor cells, mast cells, and is also found in a wide range of non-haemopoietic cell types (including melanocytes, germ cells, astrocytes, renal tubules, breast glandular epithelial cells, sweat glands, and interstitial cells of Cajal) (PMID: 10582338; 23073628). CD117 plays an important role in early hemopoiesis. It is also involved in pigmentation, fertility, gut movement, and some aspects of the nervous system (PMID: 23073628).

Notable Publications

Author	Pubmed ID	Journal	Application
Yuelong Jiang	36355485	Pharmaceuticals (Basel)	WB
Xiu-Bo Sang	27748937	Oncol Rep	
Wenbo Liu	36355419	eLife	

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Mo7e cells were subjected to SDS PAGE followed by western blot with 18696-1-AP (CD117/c-kit antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded stromal tumor tissue slide using 18696-1-AP (CD117/c-kit antibody at dilution of 1:200 (under 10x lens).



Immunohistochemical analysis of paraffin-embedded stromal tumor tissue slide using 18696-1-AP (CD117/c-kit antibody at dilution of 1:200 (under 40x lens).



Immunofluorescent analysis of (-20°C Ethanol) fixed MCF-7 cells using 18696-1-AP (CD117/c-kit antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).