## For Research Use Only

## EGFR-Specific Polyclonal antibody

Catalog Number: 18986-1-AP

Featured Product

171 Publications



**Basic Information** 

Catalog Number: GenBank Accession Number: 18986-1-AP NM\_005228

Concentration: GeneID (NCBI): 1956

Source: UNIPROT ID: Rabbit P00533

Isotype: Full Name:

epidermal growth factor receptor (erythroblastic leukemia viral (v-erbb) oncogene homolog, avian)

Calculated MW: 134 kDa Observed MW: 145-165 kDa Purification Method: Antigen affinity purification Recommended Dilutions: WB: 1:1000-1:6000

IP: 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

**Applications** 

Tested Applications: WB, IP, ELISA Cited Applications: WB, IHC, IF, IP, CoIP, ChIP Species Specificity:

human Cited Species:

human, mouse, rat, pig, sheep, goat, sus scrofa (pig)

Positive Controls

WB: HepG2 cells, MCF-7 cells, A431 cells, HeLa/RPE1 cells, NCI-H1299 cells, HeLa cells, A549 cells, LO2 cells,

NCL-H1299 cells IP: MCF-7 cells,

**Background Information** 

ERBB1, also named as EGFR, is a receptor for EGF, but also for other members of the EGF family, as TGF-alpha, amphiregulin, betacellulin, heparin-binding EGF-like growth factor, GP30 and vaccinia virus growth factor. ERBB1 is involved in the control of cell growth and differentiation. Phosphorylates MUC1 in breast cancer cells and increases the interaction of MUC1 with C-SRC and CTNNB1/beta-catenin. Isoform 2/truncated isoform of ERBB1 may act as an antagonist. EGFR is associated with lung cancer. The antibody is specific to isoform1.

## **Notable Publications**

Author	Pubmed ID	Journal	Application
Haile Zhao	34658851	Front Pharmacol	WB,IP
Taiwei Wang	36169181	Oncol Rep	WB
Jing Cai	31826234	Carcinogenesis	WB,IHC,IF

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


## Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 18986-1-AP (EGFR-Specific antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



WB result of EGFR-Specific antibody (18986-1-AP; 1:3000; incubated at room temperature for 1.5 hours) with sh-Control and sh-EGFR-Specific transfected HepG2 cells.



WB result ofanti-EGFR (18986-1-AP) in Hela and REP1 cell by Dr.Kodani, Andrew.



IP result of anti-EGFR-Specific (IP:18986-1-AP, 3ug; Detection:18986-1-AP 1:500) with MCF-7 cells lysate 2500ug.