For Research Use Only

DRD5 Polyclonal antibody

Catalog Number: 20310-1-AP

Featured Product

11 Publications



Basic Information

Catalog Number:
20310-1-AP
BC009748

Concentration:
GeneID (NCBI):
600 ug/ml
1816

Source:
UNIPROT ID:
Rabbit
P21918

Isotype:
Full Name:

IgG dopamine receptor D5
Immunogen Catalog Number: Calculated MW:

Observed MW:

50-53 kDa

477 aa, 53 kDa

Purification Method: Antigen affinity purification

Recommended Dilutions: WB: 1:500-1:3000

IHC: 1:50-1:500

FC (Intra): 0.80 ug per 10^6 cells in a

100 µl suspension

Applications

Tested Applications: WB, IHC, FC (Intra), ELISA Cited Applications:

AG14145

WB, IHC, IF Species Specificity: human, mouse, rat Cited Species: human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: mouse brain tissue, rat brain tissue

IHC : mouse brain tissue,
FC (Intra) : SH-SY5Y cells,

Background Information

DRD5, is a G-protein coupled receptor and involved in diversified physiological and pathological processes. this receptor stimulates adenylyl cyclase activity and is expressed in the limbic area of the mammalian brain(PMID:14699430). Studies have shown that DRD5 is highly expressed in colonic macrophages and DRD5 deficiency exacerbates DSS(dextran sodium sulfate)-induced colitis(PMID:34001860). in addition, DRD5 agonists were potent inhibitors of pituitary tumor growth(PMID:28613975).

Notable Publications

Author	Pubmed ID	Journal	Application
Gongliang Zhang	31491076	ACS Chem Neurosci	WB
Magdalena Kurnik-Łucka	34884737	Int J Mol Sci	IHC
Lu Liu	34001860	Cell Death Dis	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 20310-1-AP (DRD5 antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 20310-1-AP (DRD5 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 20310-1-AP (DRD5 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



mouse brain tissue were subjected to SDS PAGE followed by western blot with 20310-1-AP (DRD5 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



1x10^6 SH-SY5Y cells were intracellularly stained with 0.8 ug DRD5 Polyclonal antibody (20310-1-AP) and Coralite@488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2)(red), or 0.8 ug Rabbit IgG control Rabbit PolyAb (30000-0-AP) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).