For Research Use Only

AKT3 Polyclonal antibody

Catalog Number: 21641-1-AP 14 Publications



Purification Method:

WB: 1:500-1:1000 IHC: 1:50-1:500

Antigen affinity purification

Recommended Dilutions:

Basic Information

Catalog Number:

21641-1-AP

BC121154

Concentration:

760 µg/ml

Source:

Rabbit

Q9Y243

Isotype:

GenBank Accession Number:

BC121154

GeneID (NCBI):

10000

UNIPROT ID:

Q9Y243

Full Name:

IgG v-akt murine thymoma viral oncogene homolog 3 (protein kinase

AG16298 B, gamma)

Calculated MW:
465 aa, 54 kDa

Observed MW:
60 kDa

Applications

Tested Applications: WB, IHC, ELISA Cited Applications: WB. IHC

Species Specificity: human, mouse, rat Cited Species: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: mouse brain tissue, rat brain tissue

IHC: human tonsillitis tissue,

Background Information

AKT3, also named as PKBG, is one of 3 closely related serine/threonine-protein kinases (AKT1, AKT2 and AKT3) called the AKT kinase, and which regulate many processes including metabolism, proliferation, cell survival, growth and angiogenesis. AKT3 is a key modulator of several tumors like melanoma, glioma and ovarian cancer. Active AKT3 increases progressively during melanoma tumor progression with highest levels present in advanced-stage metastatic melanomas.

Notable Publications

Author	Pubmed ID	Journal	Application
Yang Yang	31515938	J Cell Mol Med	WB
Yong-Xia Wang	30186040	Cancer Cell Int	WB
Wei Sang	27329362	Cell Immunol	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 4006900926 E: Proteintech-CN@ptglab.com

W: ptgcn.cor

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 21641-1-AP (AKT3 antibody) at dilution of 1:800 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 21641-1-AP (AKT3 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).