For Research Use Only

Neuropilin 1/CD304 Monoclonal antibody



Catalog Number: 60067-1-lg

Featured Product

25 Publications

Basic Information

Catalog Number: 60067-1-lg Concentration: 1000 ug/ml Source:

Mouse
Isotype:
IgG1
Immunogen Catalog Number:
AG0931

1 103 kDa
Observed MW:
130 kDa

Applications

Tested Applications:
WB, IF/ICC, ELISA
Cited Applications:
WB, IF, IP, CoIP
Species Specificity:
human, pig, rabbit
Cited Species:

human, mouse, rabbit, xenopus

GenBank Accession Number:

BC007737
GeneID (NCBI):
8829
UNIPROT ID:

O 14786
Full Name:
neuropilin 1
Calculated MW:

Positive Controls:

WB: human placenta tissue, fetal human brain tissue, human plasma, 37°C incubated human placenta tissue, pig brain tissue, HeLa cells, rabbit heart tissue, MDA-MB-231 cells, A549 cells, pig heart tissue

Purification Method:

Protein G purification

Recommended Dilutions:

WB: 1:1000-1:6000 IF/ICC: 1:50-1:500

CloneNo.:

2H3F6

IF/ICC: SH-SY5Y cells, human embronic stem cells

Background Information

Neuropilin-1 (NRP1) is a 130-140 kDa transmembrane glycoprotein expressed by endothelial, dendritic, and regulatory T cells, as well as several other normal cell types and malignant tumor cells. NRP1 was first identified as a semaphorin (SEMA) receptor, involved in axonal guidance in embryonic development. NRP1 was also shown to act as a receptor for vascular endothelial growth factor (VEGF) and a promoter of angiogenesis through its interaction with VEGF-A165 (and other VEGFs) and the receptor tyrosine kinase (RTK) VEGF-R2. NRP1 plays versatile roles in angiogenesis, axon guidance, cell survival, migration, and invasion.

Notable Publications

Author	Pubmed ID	Journal	Application
Chunxi Liu	25204970	Mol Cell Biochem	WB
Emmanuel Laplantine	36093378	iScience	WB
Max Koppers	31746735	Elife	IF

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 60067-1-1g (Neuropilin 1 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.








Confocal immunofluorescent analysis of human embronic stem cells with 60067-1-lg at dilution of 1:100. DAPI was used to stain the cell nuclei (blue) at a concentration of 1ug/ml. This photo is the merge of PE and DAPI. (40X)





Immunofluorescent analysis of (-20°C Ethanol) fixed SH-SY5Y cells using 60067-1-1g(Neuropilin 1 antibody) at dilution of 1:100 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Mouse IgG(H+L).