## For Research Use Only

## B23/NPM1 Monoclonal antibody, PBS Only



Catalog Number: 60096-1-PBS

Featured Product

**Basic Information** 

Catalog Number: GenBank Accession Number: 60096-1-PBS BC002398 GeneID (NCBI): Concentration: 1mg/ml 4869 **UNIPROT ID:** Source:

Mouse P06748 Isotype: Full Name: lgG1 nucleophosmin (nucleolar

phosphoprotein B23, numatrin) Immunogen Catalog Number: Calculated MW:

AG7415

33 kDa Observed MW: 35-38 kDa

**Purification Method:** Protein G purification

CloneNo.: 4F12A3

**Applications** 

**Tested Applications:** 

WB, IHC, IF/ICC, Indirect ELISA

Species Specificity: human, mouse, rat

**Background Information** 

Nucleophosmin (NPM1,B23) is a putative ribosome assembly factor with a high affinity for peptides containing nuclear localization signals (NLSs). The transport of proteins across the nuclear envelope is a selective, multistep process involving several cytoplasmic factors. Proteins must be recognized as import substrates, dock at the nuclear pore complex and translocate across the nuclear envelope in an ATP-dependent fashion. Several cytosolic and nuclear proteins that are central to this process have been identified. The 38 kDa nuclear protein nucleophosmin is involved in ribosomal assembly and rRNA transport. It is an abundant protein that is highly phosphorylated by Cdc2 kinase during mitosis.

Storage


Storage:

Store at -80°C.


The product is shipped with ice packs. Upon receipt, store it immediately at -80°C Storage Buffer:

PBS only, pH7.3


## Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 60096-1-lg (B23/NPM1 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Alpha Tubulin Monoclonal antibody (HRP-66031) as loading control. This data was developed using the same antibody clone with 60096-1-PBS in a different storage buffer formulation




Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 60096-1-lg (B23 Antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60096-1-PBS in a different storage buffer formulation.




WB result of B23/NPM1 antibody (60096-1-lg; 1:10000; incubated at room temperature for 1.5 hours) with sh-Control and sh-B23/NPM1 transfected HepG2 cells. This data was developed using the same antibody clone with 60096-1-PBS in a different storage buffer formulation.






Various lysates were subjected to SDS PAGE followed by western blot with 60096-1-1g (B23/NPM1 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 60096-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using B23/NPM1 antibody (60096-1-lg, Clone: 4F12A3) at dilution of 1:800 and CoraLite® 488-Conjugated Goat Anti-Mouse IgG(H+L) (SA00013-1), CL594-phalloidin (red). This data was developed using the same antibody clone with 60096-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed MCF-7 cells using B23/NPM1 antibody (60096-1-1g, Clone: 4F12A3) at dilution of 1:800 and CoraLite® 488-Conjugated Goat Anti-Mouse IgG(H+L) (SA00013-1). This data was developed using the same antibody clone with 60096-1-PBS in a different storage buffer formulation.