For Research Use Only

AKT Monoclonal antibody, PBS Only (Detector)



Purification Method:

Protein G purification

CloneNo.:

2C5D1

Catalog Number: 60203-2-PBS

Featured Product

Basic Information

Catalog Number: 60203-2-PBS Concentration: 1mg/ml

UNIPROT ID: Source: Mouse P31749 Isotype: Full Name: lgG1 v-akt murine thymoma viral

Immunogen Catalog Number:

AG16695

Calculated MW: 56 kDa

> Observed MW: 56-62 kDa

oncogene homolog 1

BC000479

GeneID (NCBI):

GenBank Accession Number:

Applications

Background Information

Tested Applications:

WB, IHC, IF/ICC, IP, Cytometric bead array, Indirect

ELISA

Species Specificity: human, mouse, rat

Storage

 $The serine-threonine\ protein\ kinase\ AKT1\ is\ catalytically\ inactive\ in\ serum-starved\ primary\ and\ immortalized$ fibroblasts. Survival factors can suppress apoptosis in a transcription-independent manner by activating the serine/threonine kinase AKT1, which then phosphorylates and inactivates components of the apoptotic machinery. This antibody detects all the members of AKT with/without phospho-modification.


Storage:

Store at -80°C.


The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

PBS only, pH7.3


Selected Validation Data



WB result of AKT antibody (60203-2-lg; 1:5000; incubated at room temperature for 1.5 hours) with sh-Control and sh-AKT transfected A549 cells. This data was developed using the same antibody clone with 60203-2-PBS in a different storage buffer formulation.




Various lysates were subjected to SDS PAGE followed by western blot with 60203-2-lg (AKT antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 60203-2-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 60203-2-lg (AKT Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60203-2-PBS in a different storage buffer formulation.




Immunofluorescent analysis of (-20°C Methanol) fixed MCF-7 cells using AKT antibody (60203-2-lg, Clone: 2C5D1) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 60203-2-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 60203-2-lg (AKT Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 60203-2-PBS in a different storage buffer formulation.



HEK-293 cells were subjected to SDS PAGE followed by western blot with 60203-2-1g (AKT Antibody) and 66444-1-1g(AKT1-phospho-5473 Antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 60203-2-PBS in a different storage buffer formulation.



IP result of anti-AKT (IP:60203-2-Ig, 5ug; Detection:60203-2-Ig 1:1000) with mouse brain tissue lysate 4000ug. This data was developed using the same antibody clone with 60203-2-PBS in a different storage buffer formulation.



Cytometric bead array in cell lysate using MP50177-1, Phospho-AKT(Ser473) Monoclonal Matched Antibody Pair, PBS Only. Capture antibody: 66444-1-PBS. Detection antibody: 60203-2-PBS. Cell lysate: Non-treated HEK293 and Calyculin A treated HEK293(30 µ g/well). Nonrelated target OAT Monoclonal Matched Antibody Pair (MP50109-1P) was served as control.