### For Research Use Only

# CD23 Monoclonal antibody

Catalog Number:60208-1-lg 4 Publications



**Basic Information** 

Catalog Number: GenBank Accession Number: 60208-1-lg BC064417 GeneID (NCBI): Concentration: 1000 ug/ml 2208 **UNIPROT ID:** Source: Mouse P06734 Full Name: Isotype:

lgG1 Fc fragment of IgE, low affinity II, receptor for (CD23) Immunogen Catalog Number:

AG0425 Calculated MW: 321 aa, 36 kDa Observed MW:

36 kDa

**Purification Method:** Protein G purification

CloneNo.:

3H6B8

Recommended Dilutions: WB: 1:5000-1:50000 IHC: 1:500-1:2000 IF-P: 1:200-1:800

**Applications** 

**Tested Applications:** WB, IHC, IF-P, ELISA **Cited Applications:** WB, IHC, IF Species Specificity: human, mouse

**Cited Species:** human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

#### Positive Controls:

WB: THP-1 cells, DC2.4 cells, U-937 cells, Raji cells, Ramos cells, Daudi cells, Mo7e cells

IHC: human tonsillitis tissue, IF-P: human tonsillitis tissue,

## **Background Information**

CD23, also known as low affinity immunoglobulin epsilon Fc receptor, is a transmembrane glycoprotein present on  $a\ subpopulation\ of\ B\ lymphocytes\ in\ germinal\ centres,\ EBV-transformed\ B-lymphoblastoid\ cell\ lines,\ follicular$ dendritic cells, and a subpopulation of peripheral blood cells. CD23 has essential roles in the regulation of IgE production and in the differentiation of B-cells.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Kiyoko Maruyama	34719578	Biol Pharm Bull	WB,IF
Ivan K Chinn	34315806	Sci Signal	IF
Qing Wang	37525587	Clin Transl Med	IHC,IF

## Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**




Various lysates were subjected to SDS PAGE followed by western blot with 60208-1-lg (CD23 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 60208-1-1g (CD23 antibody) at dilution of 1:1000 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded human tonsillitis tissue using CD23 antibody (60208-1-lg, Clone: 3H6B8) at dilution of 1:400 and CoraLite®488-Conjugated Goat Anti-Mouse IgG(H+L) (\$A00013-1). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9 0)



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded human tonsillitis tissue using CD23 antibody (60208-1-lg, Clone: 3H6B8) at dilution of 1:400 and Coralite® 488-Conjugated Goat Anti-Mouse IgG(H+L) (SA00013-1). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).