For Research Use Only

P-selectin / CD62P Monoclonal antibody

www.ptglab.com

Catalog Number:60322-1-lg 33 Publications

Basic Information

Catalog Number: 60322-1-lg Concentration: 1295 ug/ml

Source: Mouse Isotype: lgG2b

Immunogen Catalog Number:

AG4124

Applications

Tested Applications: WB, IP, ELISA Cited Applications:

WB, IF Species Specificity:

human Cited Species: human, mouse, rat GenBank Accession Number:

BC028067 GeneID (NCBI): 6403

UNIPROT ID: P16109 Full Name:

selectin P (granule membrane protein protein lysate 140kDa, antigen CD62)

Calculated MW: 830 aa, 91 kDa

Observed MW: 140 kDa, 105-110 kDa

Positive Controls:

WB: human peripheral blood platelets, Jurkat cells

Purification Method:

CloneNo.:

3G2C7

Protein A purification

Recommended Dilutions:

IP: 0.5-4.0 ug for 1.0-3.0 mg of total

WB: 1:2000-1:10000

IP: K-562 cells,

Background Information

P-selectin, also known as GMP140 or CD62P, is a transmembrane glycoprotein that mediates the interaction of $activated\ endothelial\ cells\ or\ platelets\ with\ leukocytes.\ It\ is\ an\ adhesion\ molecule\ involved\ in\ the\ pathogenesis\ of\ activated\ endothelial\ cells\ or\ platelets\ with\ leukocytes.\ It\ is\ an\ adhesion\ molecule\ involved\ in\ the\ pathogenesis\ of\ activated\ endothelial\ cells\ or\ platelets\ with\ leukocytes.$ inflammation, thrombosis, and oncogenesis. P-selectin is stored in the alpha-granules of platelets and Weibel-Palade bodies of endothelial cells. Upon cell activation by agonists, P-selectin is transported rapidly to the cell

Notable Publications

Author	Pubmed ID	Journal	Application
Jinwei Pang	34566870	Front Neurol	WB,IF
LinXi Cheng	34489520	Commun Biol	FC
Chunyu Huang	33015023	Front Bioeng Biotechnol	WB


Storage


Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


Selected Validation Data







human peripheral blood platelets were subjected to SDS PAGE followed by western blot with 60322-1-Ig (P-selectin / CD62P antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



IP result of anti-P-selectin / CD62P (IP:60322-1-Ig, 5ug; Detection:60322-1-Ig 1:600) with K-562 cells lysate 1760ug.