### For Research Use Only

# Anti-Human CD28 (CD28.2)

Catalog Number:65099-1-lg 3 Publications



**Basic Information** 

Catalog Number: 65099-1-lg

Concentration: 100ug, 0.5 mg/ml

Source: Mouse

Isotype: IgG1, kappa

**Tested Applications:** 

Human

**Cited Species:** human

GenBank Accession Number:

BC093698 GeneID (NCBI):

940

P10747

ENSEMBL Gene ID: ENSG00000178562 UNIPROT ID:

Full Name: CD28 molecule Calculated MW: 220 aa, 25 kDa

**Purification Method:** 

Affinity purification

CloneNo.: CD28.2

Recommended Dilutions:

FC: 0.2 ug per 10^6 cells in 100  $\,\mu$  l

suspension

**Applications** 

FC

Cited Applications: IHC, Cell treatment Species Specificity:

Positive Controls:

FC: human PBMCs,

## **Background Information**

CD28 (T-cell-specific surface glycoprotein CD28), also known as T44 and Tp44, is a 44 kD disulfide-linked homodimeric type I glycoprotein (PMID: 2162180). It is a member of the immunoglobulin superfamily and is expressed on most T lineage cells, NK cell subsets, and plasma cells (PMID: 2162180, 8386518). CD28 may affect in vivo immune responses by functioning both as a cell adhesion molecule linking B and T lymphocytes and as the surface component of a novel signal transduction pathway (PMID: 2162180, 3021470). CD28 binds both CD80 and CD86 with a highly conserved motif MYPPY in the CDR3-like loop (PMID: 15696168, 7964482). CD28 is considered a major co-stimulatory molecule, inducing T lymphocyte activation and IL-2 synthesis, and preventing cell death (PMID: 1348520).

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Wenbo Zhao	36309750	J Transl Med	
Ya-Nan Pi	34852825	Cancer Cell Int	IHC
Guojun Yao	37354488	Aging (Albany NY)	Cell treatment


Storage

Store at 2-8°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.09% sodium azide, pH7.3

### Selected Validation Data



1X10^6 human PBMCs were surface co-stained with CL647 Plus Anti-Human CD3 and 0.2 ug Anti-Human CD28 (65099-1-lg, Clone:CD28.2) with CoraLite® 594-Conjugated Goat Anti-Mouse IgG(H+L) at dilution 1:1000. Cells were not fixed. Lymphocytes were gated.



1X10^6 human PBMCs were surface stained with CL647 Plus Anti-Human CD3 and 0.2 ug Anti-Human CD28 (65099-1-Ig, Clone:CD28.2) with CoraLite®594-Conjugated Goat Anti-Mouse IgG(H+L) at dilution 1:1000 or unstained. Cells were not fixed. Lymphocytes were gated.