

For Research Use Only

Anti-Human CD19 (HIB19)

Catalog Number: 65110-1-Ig



Basic Information

Catalog Number:

65110-1-Ig

Concentration:

100ug, 0.5 mg/ml

Source:

Mouse

Isotype:

IgG1, kappa

GenBank Accession Number:

BC006338

GeneID (NCBI):

930

ENSEMBL Gene ID:

ENSG00000177455

Full Name:

CD19 molecule

Calculated MW:

556 aa, 61 kDa

Purification Method:

Affinity purification

CloneNo.:

HIB19

Recommended Dilutions:

FC: 0.2 ug per 10⁶ cells in 100 μ l suspension

Applications

Tested Applications:

FC

Species Specificity:

Human

Positive Controls:

FC : human PBMCs,

Background Information

CD19 is a 95 kDa type I transmembrane glycoprotein belonging to the immunoglobulin superfamily (PMID: 2472450). It is expressed by B cells and follicular dendritic cells. CD19 is up-regulated at the step of B-lineage commitment during the differentiation of the hematopoietic stem cell, it remains on during subsequent stages of differentiation until finally down-regulated during terminal differentiation into plasma cells (PMID: 8528044). CD19 is involved in B cell development, activation and differentiation. It is the dominant component for the signaling complex on B cells that includes CD21 (CR2), CD81 (TAPA-1) and CD225 and acts as a critical co-receptor for BCR signal transduction (PMID: 23210908).

Storage

Storage:

Store at 2-8°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.09% sodium azide, pH7.3

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1X10⁶ human PBMCs were surface stained with 0.2 ug Anti-Human CD19 (65110-1-Ig, Clone:H1B19) or Mouse IgG1 Isotype Control and CoraLite®594-Conjugated Goat Anti-Mouse IgG(H+L) at dilution 1:500, then CL405 Anti-Human CD3. Cells were not fixed. Lymphocytes were gated.



1X10⁶ human PBMCs were surface stained with 0.2 ug Anti-Human CD19 (65110-1-Ig, Clone:H1B19) or Mouse IgG1 Isotype Control and CoraLite®594-Conjugated Goat Anti-Mouse IgG(H+L) at dilution 1:500. Cells were not fixed. Lymphocytes were gated.