For Research Use Only

Anti-Human CD3 (UCHT1)

Catalog Number: 65151-1-lg



Purification Method:

Affinity purification

Recommended Dilutions:

FC: 0.2 ug per 10^6 cells in 100 $\,\mu$ l

CloneNo.: UCHT1

suspension

Basic Information

Catalog Number: 65151-1-lg

Concentration: 500ug, 0.5 mg/ml

Mouse Isotype: IgG1, kappa

Source:

Species Specificity:

Tested Applications:

Human

GenBank Accession Number:

BC049847 GeneID (NCBI): 916

Calculated MW:

ENSEMBL Gene ID: ENSG00000198851 UNIPROT ID:

P07766 Full Name:

CD3e molecule, epsilon (CD3-TCR complex)

207 aa, 23 kDa

Positive Controls:

FC : human PBMCs,

Background Information

CD3 is a multimeric protein associated with the T-cell receptor (TCR) to form a complex involved in antigen recognition and signal transduction (PMID: 15885124). CD3 is composed of CD3 γ , δ , ϵ , and ζ chains (PMID: 1826255). It is expressed by thymocytes in a developmentally regulated manner, T cells, and some NK cells (PMID: 3289580). The TCR recognizes antigens bound to major histocompatibility complex (MHC) molecules. TCR-mediated peptide-MHC recognition is transmitted to the CD3 complex, leading to the intracellular signal transduction (PMID: 11985657).

Storage

Applications


Storage:

Store at 2-8°C. Stable for one year after shipment.


Storage Buffer

PBS with 0.09% sodium azide, pH7.3

Selected Validation Data



1X10^6 human PBMCs were surface co-stained with FITC Anti-Human CD19 and 0.2 ug Anti-Human CD3 (65151-1-lg, Clone:UCHT1) labeled with FlexAble CoraLite® Plus 647 Antibody Labeling Kit for Mouse IgG1 or Mouse IgG1 lsotype Control. Cells were not fixed. Lymphocytes were gated.



1X10^6 human PBMCs were surface stained with 0.2 ug Anti-Human CD3 (65151-1-lg, Clone:UCHT1) labeled with FlexAble CoraLite® Plus 647 Antibody Labeling Kit for Mouse IgG1 or Mouse IgG1 lsotype Control. Cells were not fixed. Lymphocytes were gated.