For Research Use Only

BMI1 Monoclonal antibody

Catalog Number:66161-1-lg

Featured Product

15 Publications

BC011652

GeneID (NCBI):

UNIPROT ID:

Full Name:

Calculated MW:

P35226

37 kDa

GenBank Accession Number:

BMI1 polycomb ring finger oncogene



Basic Information

Applications

Catalog Number: 66161-1-lg Concentration:

1000 ug/ml Source: Mouse Isotype: lgG2b

Immunogen Catalog Number:

AG21284

Observed MW:

37-43 kDa

Tested Applications:

WB, IHC, IF/ICC, IP, ELISA

Cited Applications:

WB, IHC, IF

Species Specificity: human, mouse, monkey

Cited Species:

human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: PC-12 cells, HT-1080 cells, U2OS cells, U-937 cells

Purification Method:

Protein A purification

Recommended Dilutions:

IP: 0.5-4.0 ug for 1.0-3.0 mg of total

WB: 1:1000-1:4000

protein lysate

IHC: 1:50-1:500

IF/ICC: 1:500-1:2000

CloneNo.:

2E10A11

IP: HEK-293 cells,

IHC: human breast cancer tissue, human colon cancer,

human lymphoma tissue

IF/ICC: HEK-293 cells, COS-7 cells

Background Information

BMI-1 is one of polycomb group genes, which together with Ring1 strongly enhances the E3 ubiquitin ligase activity of the Ring2 catalytic subunit. Bmi 1 plays an important role in the regulation of cell proliferation and senescence through repression of the p16Ink4a and p19Arf genes and is required for maintenance of adult hematopoietic and neural stem cells. The antibody reacts with the 37kd BMI1 protein.

Notable Publications

Author	Pubmed ID	Journal	Application
Anindya Dey	29158468	Mol Cancer Ther	WB,IHC
Guan-Zhong Qiu	29788550	Cancer Sci	WB,IF
Mingzhu Tan	34391775	Exp Cell Res	WB,IF

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




PC-12 cells were subjected to SDS PAGE followed by western blot with 66161-1-1g (BMI1 antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.



IP result of anti-BMI1 (IP:66161-1-Ig, 3ug; Detection:66161-1-Ig 1:1000) with HEK-293 cells lysate 1800ug.



Immunohistochemical analysis of paraffinembedded human lymphoma tissue slide using 66161-1-lg (BMI1 Antibody) at dilution of 1:50 under 40x lens.



Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 66161-1-Ig (BMI1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HEK-293 cells using BMI1 antibody (66161-1-1g, Clone: 2E10A11) at dilution of 1:1000 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).