For Research Use Only

CD13 Monoclonal antibody, PBS Only (Capture)

proteintech
Antibodies | ELISA kits | Proteins
www.ptglab.com

Purification Method:

CloneNo.:

2D8D11

Protein A purification

Catalog Number: 66211-1-PBS

Basic Information

Catalog Number:

66211-1-PBS

Size: GeneID (NCBI):

1mg/ml 29 Source: EN

Source: ENSEMBL Gene ID:
Mouse ENSG00000166825
Isotype: UNIPROT ID:
IgG2a P15144
Immunogen Catalog Number: Full Name:

AG5976 alanyl (membrane) aminopeptidase

Calculated MW: 110 kDa Observed MW: 150 kDa

GenBank Accession Number:

BC058928

Applications

Tested Applications:

WB, IHC, IF/ICC, IF-P, Cytometric bead array, Indirect

ELISA

Species Specificity:

human, mouse, rat

Background Information

CD13, also named as APN, ANPEP(aminopeptidase N) or PEPN, is belongs to the peptidase M-1 family. CD13 is a heavily glycosylated, ~150-240 kDa, type-II membrane, expressed by most cells of myeloid origin including monocytes, macrophages, granulocytes, and their hematopoietic precursors. It is also abundantly expressed in the brush border of epithelial cells from renal proximal tubules and small intestine, in prostatic epithelial cells, in bile duct canaliculi, in mast cells, and, in some cases, in fibroblasts and smooth muscle cells. CD13 is a multifunctional protein and plays varying roles in cell migration, cell proliferation, cell differentiation and so on. CD13 participates in angiogenesis generating and modulating angiogenic signals, and can be a marker of angiogenic vessels. CD13 is also a pan-myeloid marker, present on mature granulocytes and monocytes. (PMID: 8805662, 10098327, 18603472, 18097955, 17897790, 17888402, 21339174)

Storage


Storage:

Store at -80°C.


The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer: PBS Only


Selected Validation Data




Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 66211-1-lg (CD13 antibody) at dilution of 1:10000 (under 4x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66211-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded human kidney tissue slide using 66211-1-Ig (CD13 antibody) at dilution of 1:10000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66211-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 66211-1-lg (CD13 antibody) at dilution of 1:10000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66211-1-PBS in a different storage buffer formulation.




Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded mouse kidney tissue using CD13 antibody (66211-1-lg, Clone: 2D8D11) at dilution of 1:400 and Coralite® 488-Conjugated Goat Anti-Mouse IgG(H+L) (SA00013-1). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66211-1-PBS in a different storage buffer formulation.




Immunofluorescent analysis of (4% PFA) fixed THP-1 cells using CD13 antibody (66211-1-lg, Clone: 2D8D11) at dilution of 1:800 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 66211-1-PBS in a different storage buffer formulation.




U-937 cells were subjected to SDS PAGE followed by western blot with 66211-1-1g (CD13 antibody at dilution of 1:4000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66211-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 66211-1-lg (CD13 antibody) at dilution of 1:10000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66211-1-PBS in a different storage buffer formulation.




Immunofluorescent analysis of (4% PFA) fixed human kidney tissue using 66211-1-1g (CD13 antibody) at dilution of 1:100 and Alexa Fluor 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 66211-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human kidney tissue slide using 66211-1-Ig (CD13 antibody) at dilution of 1:10000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66211-1-PBS in a different storage buffer formulation.

Immunohistochemical analysis of paraffinembedded human kidney tissue slide using 66211-1-Ig (CD13 antibody) at dilution of 1:10000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66211-1-PBS in a different storage buffer formulation.



Cytometric bead array standard curve of MP50521-1, ANPEP Monoclonal Matched Antibody Pair, PBS Only. Capture antibody: 66211-1-PBS. Detection antibody: 66211-2-PBS. Standard:Ag5976. Range: 0.781-100 ng/mL