## For Research Use Only

## PD-L1/CD274 Monoclonal antibody, PBS Only

Proteintech®
Antibodies | ELISA kits | Proteins
www.ptglab.com

**Purification Method:** 

Protein G purification

CloneNo.:

2B11D11

Catalog Number: 66248-1-PBS

**Featured Product** 

**Basic Information** 

Catalog Number: 66248-1-PBS Concentration: 1000 ug/ml

Source:
Mouse
Isotype:
IgG1
Immunogen Catalog Number:

AG12443 290 aa, 33 kDa Observed MW:

Observed MW: 45-50 kDa, 33 kDa

CD274 molecule Calculated MW:

GenBank Accession Number:

BC074984

29126

Q9NZQ7
Full Name:

GeneID (NCBI):

**UNIPROT ID:** 

**Applications** 

Tested Applications: WB, IHC, IF/ICC, IF-P, ELISA Species Specificity: human, mouse, rat, pig

## **Background Information**

Programmed cell death ligand 1 (PD-L1, CD274, or B7-H1), is the first member of B7 family to be discovered. B7 family molecules are type I transmembrane proteins belonging to the immunoglobulin superfamily. In concert with their CD28 family receptors, the B7s are key regulators of the adaptive immune response. PD-L1 is suggested as a negative regulator of T and B cell, and plays important role in mediating tolerance of lymphocytes to self-antigens. It is also involved in the costimulatory signal, essential for T-cell proliferation and production of IL10 and IFNG, in an IL2-dependent and a PD-1-independent manner. PD-L1 is a 290 aa transmembrane protein with a calculated molecular weight of 33 kDa, it is predicted to be 27-30 kDa after signal peptide cleavage (PMID: 25609200; 17076679). The apparent molecular weight has also been reported as 45-70 kDa, major glycosylated form of 45-50 kDa and multiple post-translational modifications form of 65-70 kDa (PMID: 18760278; 16493058).

Storage


Storage:

Store at -80°C.


The product is shipped with ice packs. Upon receipt, store it immediately at -80°C Storage Ruffer.

PBS only, pH7.3

## Selected Validation Data



human placenta tissue were subjected to SDS PAGE followed by western blot with 66248-1-lg (PD-L1/CD274 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66248-1-PBS in a different storage buffer formulation.




Untreated and IFN gamma treated A375 cells and A549 cells were subjected to SDS PAGE followed by western blot with 66248-1-1g (PD-L1/CD274 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66248-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 66248-1-1g (PD-L1/CD274 antibody) at dilution of 1:10000 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66248-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 66248-1-lg (PD-L1/CD274 antibody) at dilution of 1:10000 (under 4x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66248-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (10% Formaldehyde) fixed HeLa cells using 66248-1-Ig(PD-L1/CD274 antibody) at dilution of 1:300 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 66248-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded human placenta tissue using PD-L1/CD274 antibody (66248-1-lg, Clone: 2B11D11) at dilution of 1:800 and Coralite® 488-Conjugated Goat Anti-Mouse IgG(H+L) (SA00013-1). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66248-1-PBS in a different storage buffer formulation.