#### For Research Use Only

# TAU Monoclonal antibody

Catalog Number:66499-1-lg Featured Product

24 Publications



**Basic Information** 

Catalog Number: GenBank Accession Number: 66499-1-lg BC000558 GeneID (NCBI): Concentration: 1000 ug/ml 4137

**UNIPROT ID:** Source: Mouse P10636 Full Name: Isotype:

IgG2c microtubule-associated protein tau

Immunogen Catalog Number: Calculated MW: AG21926 37-46, 79-81 kDa

> Observed MW: 100 kDa

**Applications** 

**Tested Applications:** WB, IHC, ELISA Cited Applications: WB, IHC, IF Species Specificity:

human, mouse, rat, pig

**Cited Species:** 

human, mouse, rat, rabbit

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

**Positive Controls:** 

WB: HeLa cells, Y79 cells, pig brain tissue, SH-SY5Y cells, U-251 cells, Neuro-2a cells, rat brain tissue, mouse brain tissue

**Purification Method:** 

Protein A purification

Recommended Dilutions:

WB: 1:1000-1:30000 IHC: 1:200-1:800

CloneNo.:

1E9A8

IHC: human gliomas tissue, mouse brain tissue

## **Background Information**

The microtubule-associated protein TAU (MAPT or TAU) is encoded by MAPT gene, which locates on human chromosome 17q21, binds to the tubulin subunit of microtubule and promotes its assembly and stability. Most TAU is expressed in neurons, and TAU isoform is expressed in the peripheral nervous system while the others are expressed in the central nervous system. TAU links axonal microtubules with C-terminus to neural plasma  $membrane\ components\ with\ its\ N-terminus,\ suggesting\ the\ participation\ in\ intracellular\ signal\ transduction\ and$ neuron's development and viability. Various isoforms of Tau exist due to the alternative splicing, and short isoforms around 45-69 kDa and long isoforms around 100-110 kDa have been reported in different literature (PMID:8752131,15965697, 12485403). Present monoclonal anti-Tau antibody can detect approx 100-kDa bands in brain tissues.

## Notable Publications

| Author                    | Pubmed ID | Journal              | Application |
|---------------------------|-----------|----------------------|-------------|
| Estibaliz Santiago-Mujika | 36606207  | J Alzheimers Dis Rep | WB          |
| Jiqu Xu                   | 31050371  | J Pineal Res         | WB          |
| Nicholas E. Albrecht      | 35880013  | Cell Rep Methods     | IF          |

Storage

Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol, pH7.3


Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## **Selected Validation Data**




WB result of TAU antibody (66499-1-Ig; 1:30000; incubated at room temperature for 1.5 hours) with sh-Control and sh-TAU transfected HeLa cells.



Various lysates were subjected to SDS PAGE followed by western blot with 66499-1-1g (TAU antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 66499-1-1g (TAU antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human gliomas tissue slide using 66499-1-Ig (TAU antibody) at dilution of 1:400 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human gliomas tissue slide using 66499-1-Ig (TAU antibody) at dilution of 1:400 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).