For Research Use Only

STAT1 Monoclonal antibody, PBS Only

Catalog Number:66545-1-PBS Featured Product



Purification Method:

Protein A purification

CloneNo.:

1F7C6

Basic Information

Catalog Number: 66545-1-PBS Concentration:

1000 $\,\mu$ g/ml Source: Mouse Isotype: IgG2a

Immunogen Catalog Number:

AG0199

GeneID (NCBI): 6772 **UNIPROT ID:**

> Full Name: signal transducer and activator of

GenBank Accession Number:

BC002704

P42224

84 kDa

Calculated MW: 83 kDa Observed MW:

transcription 1, 91kDa

Applications

Tested Applications:

WB, IHC, IF/ICC, Indirect ELISA

Species Specificity: human, mouse, rat

Background Information

STAT1 (signal transducers and activators of transcription 1) is a member of the STAT protein family. In response to $cytokines\ and\ growth\ factors, STAT\ family\ members\ are\ phosphory lated\ by\ the\ receptor\ associated\ kinases,\ and$ then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. Various ligands, including interferons- and -, EGF, PDGF and IL-6, can activate STAT1. STAT1 protein mediates the expression of a variety of genes, which is thought to be important for cell viability in response to different cell stimuli and pathogens.

Storage


Storage:

Store at -80°C.


The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer: PBS only, pH7.3


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 66545-1-lg (STAT1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Alpha Tubulin Monoclonal antibody (HRP-66031) as loading control. This data was developed using the same antibody clone with 66545-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human ovary tumor tissue slide using 66545-1-lg (STAT1 antibody) at dilution of 1:400 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66545-1-PBS in a different storage buffer formulation.



WB result of STAT1 antibody (66545-1-lg; 1:8000; incubated at room temperature for 1.5 hours) with sh-Control and sh-STAT1 transfected HEK-293 cells. This data was developed using the same antibody clone with 66545-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed IFN alpha treated HeLa cells using STAT1 antibody (66545-1-Ig, Clone: 1F7C6) at dilution of 1:800 and Coralite® 488-Conjugated Goat Anti-Mouse IgG(H+L) (SA00013-1), CL594-Phalloidin (red). This data was developed using the same antibody clone with 66545-1-PBS in a different storage buffer formulation.