

For Research Use Only

KU70,XRCC6 Monoclonal antibody, PBS Only

Catalog Number: 66607-1-PBS



Basic Information

Catalog Number: 66607-1-PBS	GenBank Accession Number: BC008343	Purification Method: Protein A purification
Concentration: 1mg/ml	GeneID (NCBI): 2547	CloneNo.: 3C12D9
Source: Mouse	UNIPROT ID: P12956	
Isotype: IgG1	Full Name: X-ray repair complementing defective repair in Chinese hamster cells 6	
Immunogen Catalog Number: AG21301	Calculated MW: 75 kDa	
	Observed MW: 70 kDa	

Applications

Tested Applications:
WB, IHC, IF/ICC, IF-P, IF-Fro, FC (Intra), Indirect ELISA

Species Specificity:
human, mouse, rat

Background Information

X-ray repair cross-complementing protein 6(XRCC6)gene encodes subunit p70 of the p70/p80 autoantigen which has a role in DNA repair or transposition. DNA helicase II complex binds preferentially to fork-like ends of double-stranded DNA in a cell cycle-dependent manner, which may be mediated by XRCC6. XRCC6/5 can form a heterodimer, which function as a regulatory subunit of the DNA-dependent protein kinase complex DNA-PK by increasing the affinity of the catalytic subunit PRKDC to DNA, and also it can stabilize broken DNA ends and bringing them together. Otherwise, XRCC5/6 heterodimer together with APEX1 can negatively regulate of transcription.

Storage

Storage:
Store at -80°C.
The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer:
PBS only, pH7.3

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 66607-1-Ig (KU70, XRCC6 antibody) at dilution of 1:50000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control. This data was developed using the same antibody clone with 66607-1-PBS in a different storage buffer formulation.




Various lysates were subjected to SDS PAGE followed by western blot with 66607-1-Ig (KU70, XRCC6 antibody) at dilution of 1:50000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66607-1-PBS in a different storage buffer formulation.




Immunofluorescent analysis of (4% PFA) fixed human breast cancer tissue using 66607-1-Ig (KU70, XRCC6 antibody) at dilution of 1:100 and CoraLite488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 66607-1-PBS in a different storage buffer formulation.




Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using KU70, XRCC6 antibody (66607-1-Ig, Clone: 3C12D9) at dilution of 1:800 and CoraLite488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 66607-1-PBS in a different storage buffer formulation.




Immunofluorescent analysis of (4% PFA) fixed HeLa cells using KU70, XRCC6 antibody (66607-1-Ig, Clone: 3C12D9) at dilution of 1:2000 and CoraLite488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L), CL594-Phalloidin (red). This data was developed using the same antibody clone with 66607-1-PBS in a different storage buffer formulation.



1X10⁶ HepG2 cells were intracellularly stained with 0.4 ug Anti-Human KU70, XRCC6 (66607-1-Ig, Clone: 3C12D9) and CoraLite488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Mouse IgG1 Isotype Control (MOPC-21) (65124-1-Ig, Clone: MOPC-21) (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011). This data was developed using the same antibody clone with 66607-1-PBS in a



Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 66607-1-Ig (KU70, XRCC6 antibody) at dilution of 1:4000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66607-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed frozen OCT-embedded mouse breast cancer using KU70, XRCC6 antibody (66607-1-Ig, Clone: 3C12D9) at dilution of 1:400 and CoraLite488-Conjugated Goat Anti-Mouse IgG(H+L) (SA00013-1). This data was developed using the same antibody clone with 66607-1-PBS in a different storage buffer formulation.