

For Research Use Only

# EIF2AK2, PKR Monoclonal antibody

Catalog Number: 66646-1-Ig **1 Publications**



## Basic Information

Catalog Number: 66646-1-Ig	GenBank Accession Number: BC093676	Purification Method: Protein G purification
Concentration: 1000 ug/ml	GeneID (NCBI): 5610	CloneNo.: 2A8A4
Source: Mouse	UNIPROT ID: P19525	Recommended Dilutions: WB: 1:2000-1:10000 IHC: 1:50-1:500
Isotype: IgG1	Full Name: eukaryotic translation initiation factor 2-alpha kinase 2	
Immunogen Catalog Number: AG13010	Calculated MW: 551 aa, 62 kDa	
	Observed MW: 65-68 kDa	

## Applications

Tested Applications: WB, IHC, ELISA	Positive Controls:
Cited Applications: WB	WB : HeLa cells, HEK-293 cells, MCF-7 cells, Jurkat cells
Species Specificity: human	IHC : human testis tissue, human kidney tissue
Cited Species: human	
<b>Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0</b>	

## Background Information

The eukaryotic translation initiation factor 2-alpha kinase 2 (EIF2AK2, also known as PKR or PRKR) is an important component of the host innate immune antiviral response. The product of the HSAEIF2AK2 gene is a 68 kDa protein that is ubiquitously expressed at low levels. The N-terminal part of this protein contains two dsRNA binding motifs (dsRBM). The C-terminal EIF2S1 kinase domain includes 12 sub-domains that are conserved in several Species (PMID:16954046). This antibody is specific to EIF2AK2.

## Notable Publications

Author	Pubmed ID	Journal	Application
Tomoya Kitano	38761636	Biochem Biophys Res Commun	WB

## Storage

Storage:  
Store at -20°C. Stable for one year after shipment.  
Storage Buffer:  
PBS with 0.02% sodium azide and 50% glycerol, pH7.3  
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 66646-1-Ig (EIF2AK2 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human testis tissue slide using 66646-1-Ig (EIF2AK2 antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human testis tissue slide using 66646-1-Ig (EIF2AK2 antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).