### For Research Use Only

# Calnexin Monoclonal antibody

Catalog Number:66903-1-lg Featured Product

79 Publications

BC003552

GeneID (NCBI):

**UNIPROT ID:** 

Full Name:

calnexin Calculated MW:

P27824

GenBank Accession Number:



**Basic Information** 

Catalog Number: 66903-1-lg Concentration:

1000 ug/ml Source:

Mouse

Isotype: lgG1 Immunogen Catalog Number:

AG25535

90 kDa Observed MW: 90 kDa

**Purification Method:** 

Protein G purification

CloneNo.: 2A2C6

Recommended Dilutions:

WB: 1:5000-1:20000 IHC: 1:1000-1:4000

**Applications** 

**Tested Applications:** WB, IHC, ELISA Cited Applications: WB, IHC, IF Species Specificity: human, mouse

**Cited Species:** human, mouse, rat, pig, rabbit

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

**Positive Controls:** 

WB: HeLa cells, HEK-293 cells, HepG2 cells, Jurkat cells, K-562 cells, HSC-T6 cells, PC-12 cells, NIH/3T3

cells, RAW 264.7 cells

IHC: human hepatocirrhosis tissue,

# **Background Information**

Calnexin is a molecular chaperone that resides in the endoplasmic reticulum (ER) and participates in the folding and assembly of newly synthesized proteins. Calnexin is highly abundant in ER and is frequently used as an ER

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Pu Song	36125406	EMBO Rep	IF
Folnetti A Alvarez	36114323	Breast Cancer Res Treat	WB
Xueqiang Peng	32949647	Biochim Biophys Acta Mol Cell Res	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**




Various lysates were subjected to SDS PAGE followed by western blot with 66903-1-1g (Calnexin antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.




WB result of Calnexin antibody (66903-1-Ig; 1:10000; incubated at room temperature for 1.5 hours) with sh-Control and sh-Calnexin transfected HEK-293 cells.



HEK-293 cells and HEK-293-derived exosomes (HEK-293-exo) were subjected to SDS PAGE followed by western blot with 66903-1-lg (Calnexin antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human hepatocirrhosis tissue slide using 66903-1-Ig (Calnexin antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human hepatocirrhosis tissue slide using 66903-1-Ig (Calnexin antibody) at dilution of 1:2000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).