

For Research Use Only

CDC25C Monoclonal antibody

Catalog Number: 66912-1-Ig 6 Publications



Basic Information

Catalog Number: 66912-1-Ig	GenBank Accession Number: BC019089	Purification Method: Protein G purification
Concentration: 1000 ug/ml	GeneID (NCBI): 995	CloneNo.: 2B6B12
Source: Mouse	UNIPROT ID: P30307	Recommended Dilutions: WB 1:5000-1:50000
Isotype: IgG1	Full Name: cell division cycle 25 homolog C (S. pombe)	
Immunogen Catalog Number: AG10240	Calculated MW: 473 aa, 53 kDa	
	Observed MW: 46-53 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : HeLa cells, HepG2 cells, HEK-293 cells, Jurkat cells, K-562 cells, HSC-T6 cells, PC-12 cells, NIH/3T3 cells, Raji cells
Cited Applications: WB	
Species Specificity: human, mouse, rat	
Cited Species: human, mouse	

Background Information

The cell division cycle 25 (CDC25) family of proteins are highly conserved dual specificity phosphatases that activate cyclin-dependent kinase (CDK) complexes, which in turn regulate progression through the cell division cycle. CDC25C (cell division cycle 25 homolog C) is one of the 3 isoforms of CDC25 in mammalian cells. It is present in the cytoplasm of asynchronously growing human cells (PMID:10330186). And it may be phosphorylated by its own substrate, an active cdc2-cyclin B complex, to create an autoactivation loop (PMID:7937793). It can be hyperphosphorylated to get the active form of 70 kDa (PMID:18296513). CDC25C has some isoforms with molecular mass of 46-53 kDa.

Notable Publications

Author	Pubmed ID	Journal	Application
Silu Hou	36427601	Food Chem Toxicol	WB
Jun Ma	34820015	Oncol Lett	WB
Ming Li	36526012	Toxicology	WB


Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol, pH7.3
Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:
T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 66912-1-Ig (CDC25C antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control.



Various lysates were subjected to SDS PAGE followed by western blot with 66912-1-Ig (CDC25C antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.