For Research Use Only

SARS-CoV-2 Nucleocapsid Phosphoprotein Monoclonal antibody



Catalog Number:67666-1-lg 3 Publications

Basic Information

Catalog Number:

GenBank Accession Number:

Purification Method:

67666-1-lg

NC_045512 GeneID (NCBI): Protein G purification

Concentration: 1000 ug/ml

43740575

CloneNo.:

Source:

Full Name:

1B3C3

Mouse

COVID-19 N Protein

Recommended Dilutions: WB 1:5000-1:50000

Isotype: lgG1

Immunogen Catalog Number:

AG30676

Applications

Tested Applications:

WB, ELISA

Positive Controls: WB: Ag30676,

Species Specificity:

ELISA: Recombinant protein,

virus **Cited Species:**

mouse

Background Information

The nucleocapsid (N) protein has multiple functions including formation of nucleocapsids, signal transduction virus budding, RNA replication, and mRNA transcription. N protein is an important antigen for coronavirus, and it is normally highly conserved, with a molecular weight of about 50 kDa. it can be used as a marker in diagnostic assays due to its high immunogenicity (PMID: 32416961, PMID: 32235387).67666-1-lg can be used as capture antibody. 67666-2-Ig can be used as detection antibody.

Notable Publications

Author	Pubmed ID	Journal	Application
Marina Pribanić Matešić	35216036	Viruses	
I Novodchuk	35512584	Biosens Bioelectron	
Zhaohuan Wang	39287388	J Virol	

Storage


Storage:

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



Indirect ELISA was carried out by coating eukaryotic expressed N protein at 70 ng/well followed by blocking and adding serial diluted primary antibody 67666-1-lg and 67666-2-lg respectively. Signal was developed with TMB and stopped by H2SO4. Signal strength was measured by absorbance at 450 nm.



Sandwich ELISA was carried out by coating 67666-1-Ig at 80 ng/well followed by blocking and adding different concentration of eukaryotic expressed N protein (0.5-1000 ng/mL). HRP-conjugated clone 67666-2-Ig was used at 1 μ g/mL for detection. Signal was developed with TMB and stopped by H2SO4 . Signal strength was measured by absorbance at 450 nm.



E.coli expressed SARS-CoV-2 Nucleocapsid Phosphoprotein (Cat.NO. Ag30676) was subjected to SDS-PAGE followed by western blot with 67666-1-Ig and 67666-2-Ig at various work concentration.