

For Research Use Only

NeutraKine® IFN-gamma Mouse McAb

Catalog Number: 69007-1-Ig **2 Publications**



Basic Information

Catalog Number:	GenBank Accession Number:	Purification Method:
69007-1-Ig	GeneID (NCBI):	Protein G purification
Concentration:	3458	CloneNo.:
Source:	ENSEMBL Gene ID:	1E10G7
Mouse	ENSG00000111537	Recommended Dilutions:
Isotype:	Full Name:	Neutralization: 1:10-1:100
IgG1	IFN gamma	ELISA: 1:10-1:100
Immunogen Catalog Number:		
HZ-1301		

Applications

Tested Applications:	Positive Controls:
Neutralization, ELISA	Neutralization: TNF alpha treated HT-29 cells, HT-29 cell
Cited Applications:	ELISA: Recombinant protein,
Neutralization, Cell treatment	
Species Specificity:	
human	
Cited Species:	
human	

Background Information

Interferon gamma (IFNG) is a soluble cytokine that is the only member of the type II class of interferons. It is secreted by Th1 cells, cytotoxic T cells and NK cells. The cytokine is associated with antiviral, immunoregulatory and anti-tumor properties and is a potent activator of macrophages. It plays crucial roles in pathogen clearance. Aberrant IFNG expression is associated with a number of autoinflammatory and autoimmune diseases. It has been identified in many studies as a biomarker for pleural tuberculosis (TB). Mutations in this gene are associated with aplastic anemia.

This antibody can be used to neutralize the bioactivity of Interferon gamma.

Notable Publications

Author	Pubmed ID	Journal	Application
Shiying Tang	40280928	Nat Commun	Cell treatment
Guanzhan Liang	38423357	Cell Mol Gastroenterol Hepatol	Neutralization

Storage

Storage:
Lyophilized antibodies are stable for 1 year from the date of receipt if stored between (-20°C) and (-80°C). Upon reconstitution we recommend that the solution can be stored at (4°C) for short term or at (-20°C) to (-80°C) for long term. Repeated freeze thaw cycles should be avoided with reconstituted products.

Storage Buffer:
Sterile PBS, pH7.4

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Recombinant human IFN-gamma (Cat.NO. HZ-1301) inhibits HT-29 cell line (human colorectal adenocarcinoma cell line) proliferation in the prescence of 0.5 ng/mL TNF alpha in a dose-dependent manner (blue curve, refer to bottom X-left Y axis). The activity of human IFN-gamma (2ng/mL HZ-1301 axis) is neutralized by mouse anti-human IFN-gamma monoclonal antibody 69007-1-Ig at serial dose (refer to top X-right Y axis). The ND50 is typically 100-



Recombinant human IFN-gamma (Cat.NO. HZ-1301) inhibits HT-29 cell line (human colorectal adenocarcinoma cell line) proliferation in the prescence of 0.5 ng/mL TNF alpha in a dose-dependent manner (blue curve, refer to bottom X-left Y axis). The activity of human IFN-gamma (5 ng/mL HZ-1301 axis) is neutralized by mouse anti-human IFN-gamma monoclonal antibody 69007-1-Ig at serial dose (refer to top X-right Y axis). The ND50 is typically 100-



Indirect ELISA was carried out by coating recombinant Human IFN gamma (Cat.NO. HZ-1301) at 70 ng/well followed by blocking and adding serial diluted IFN gamma antibody 69007-1-Ig and 69507-1-Ig respectively. Signal was developed with TMB and stopped by H2SO4. Signal strength was measured by absorbance at 450 nm.