

For Research Use Only

APC Anti-Human CD19 (HIB19)

Catalog Number: APC-65110

1 Publications



Basic Information

Catalog Number:

APC-65110

Concentration:

100 tests, 5 µl/test

Source:

Mouse

Isotype:

IgG1, kappa

GenBank Accession Number:

BC006338

GeneID (NCBI):

930

ENSEMBL Gene ID:

ENSG00000177455

UNIPROT ID:

P15391

Full Name:

CD19 molecule

Calculated MW:

556 aa, 61 kDa

Purification Method:

Affinity purification

CloneNo.:

HIB19

Recommended Dilutions:

FC: 5 µl per 10⁶ cells in 100 µl suspension

Excitation/Emission maximum wavelengths:

650 nm / 660 nm

Applications

Tested Applications:

FC

Cited Applications:

FC

Species Specificity:

human

Cited Species:

human

Positive Controls:

FC : human PBMCs,

Background Information

CD19 is a 95 kDa type I transmembrane glycoprotein belonging to the immunoglobulin superfamily (PMID: 2472450). It is expressed by B cells and follicular dendritic cells. CD19 is up-regulated at the step of B-lineage commitment during the differentiation of the hematopoietic stem cell, it remains on during subsequent stages of differentiation until finally down-regulated during terminal differentiation into plasma cells (PMID: 8528044). CD19 is involved in B cell development, activation and differentiation. It is the dominant component for the signaling complex on B cells that includes CD21 (CR2), CD81 (TAPA-1) and CD225 and acts as a critical co-receptor for BCR signal transduction (PMID: 23210908).

Notable Publications

Author	Pubmed ID	Journal	Application
Qiuzi Long	40091553	Adv Sci (Weinh)	FC

Storage

Storage:

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:

PBS with 0.09% sodium azide and 0.5% BSA, pH7.3

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com

W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1x10⁶ human PBMCs were surface stained with 5 ul APC Anti-Human CD19 (HIB19) (APC-65110, Clone: HIB19) or APC Mouse IgG1 Isotype Control (MOPC-21) (APC-65124, Clone: MOPC-21). Cells were co-stained with Anti-Human CD3. Cells were not fixed.