For Research Use Only

APC Anti-Human CD19 (SJ25C1)

Catalog Number: APC-65145



Purification Method:

Basic Information

Catalog Number: GenBank Accession Number: APC-65145 BC006338

BC006338 Affinity purification
Genel D (NCBI): CloneNo.:

Concentration: GeneID (Notests, 5 µl/test 930

930 SJ25C1

ENSEMBL Gene ID: Recommended Dilutions:

ENSG00000177455 FC: 5 ul per 10^6 cells in 100 µl

UNIPROT ID: suspension

Isotype: UNIPROT ID: IgG1, kappa P15391

P15391 Excitation/Emission maxima
Full Name: wavelengths:
CD19 molecule 650 nm / 660 nm

Calculated MW: 556 aa, 61 kDa

Applications

Tested Applications:

FC

Species Specificity:

human

Source:

Mouse

Positive Controls:

FC: human peripheral blood lymphocytes,

Background Information

CD19 is a 95 kDa type I transmembrane glycoprotein belonging to the immunoglobulin superfamily (PMID: 2472450). It is expressed by B cells and follicular dendritic cells. CD19 is up-regulated at the step of B-lineage commitment during the differentiation of the hematopoietic stem cell, it remains on during subsequent stages of differentiation until finally down-regulated during terminal differentiation into plasma cells (PMID: 8528044). CD19 is involved in B cell development, activation and differentiation. It is the dominant component for the signaling complex on B cells that includes CD21 (CR2), CD81 (TAPA-1) and CD225 and acts as a critical co-receptor for BCR signal transduction (PMID: 23210908).

Storage


Storage

Store at 2-8 °C. Avoid exposure to light. Stable for one year after shipment.


Storage Buffer:

PBS with 0.09% sodium azide and 0.1% gelatin, pH7.2

Selected Validation Data



100 ul human peripheral blood were surface stained with FITC Anti-Human CD3 (FITC-65151, Clone: UCHT1) and 5.00 ul APC Anti-Human CD19 (APC-65145, Clone: SJ25C1). Lymphocytes were gated for analysis. Cells were not fixed.



100 ul human peripheral blood were surface stained with FITC Anti-Human CD3 (FITC-65151, Clone: UCHT1) and APC Mouse IgG1 isotype control antibody. Lymphocytes were gated for analysis. Cells were not fixed.