For Research Use Only

CoraLite® Plus 488-conjugated Perilipin 3/TIP47 Polyclonal antibody



Catalog Number:CL488-10694 Featured Product

Basic Information

Catalog Number: GenBank Accession Number: CL488-10694 BC007566 GeneID (NCBI): Concentration: 1000 ug/ml 10226 **UNIPROT ID:** Source: Rabbit 060664

Full Name: Isotype: mannose-6-phosphate receptor

binding protein 1 Immunogen Catalog Number: AG1028 Calculated MW:

47 kDa

Observed MW: 47 kDa

Purification Method:

Antigen affinity purification Recommended Dilutions:

IF/ICC: 1:50-1:500

FC (Intra): 0.40 ug per 10⁶ cells in a

100 µl suspension

Excitation/Emission maxima

wavelengths: 493 nm / 522 nm

Applications

Tested Applications: IF/ICC, FC (Intra) Species Specificity:

Positive Controls:

IF/ICC: oleic acid treated HeLa cells, oleic acid treated

HUVEC cells

FC (Intra): HeLa cells,

Background Information

Mannose 6-phosphate receptors (M6PRs) transport newly synthesized lysosomal hydrolases from the Golgi to prelysosomes and then return to the Golgi for another round of transport. M6PRBP1 (mannose-6-phosphate receptor binding protein 1), also known as TIP47, PLIN3 or PP17, interacts with the cytoplasmic domains of both cation $independent\ and\ cation-dependent\ M6PRs,\ and\ is\ required\ for\ endosome-to-Golgi\ transport.\ In\ addition\ to\ M6PR$ recycling, M6PRBP1 plays a role in lipid droplet biogenesis, and is also implicated in rhodopsin photobleaching and viral infection. M6PRBP1 has been found to be expressed in a variety of human tissues (including colon, liver and lung parenchyme, mammary gland, and skin) and is overexpressed in certain cancer cell lines. It binds to lipid droplets and also occurs in cytosol and on endosomal membranes.

Storage


Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.


PBS with 50% glycerol, 0.05% Proclin300, 0.5% BSA, pH7.3

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



Immunofluorescent analysis of (-20°C Ethanol) fixed oleic acid treated HUVEC cells using CoraLite® Plus 488 TIP47 antibody (CL488-10694) at dilution of 1:200.



Immunofluorescent analysis of (-20°C Ethanol) fixed oleic acid treated HeLa cells using CoraLite® Plus 488 TIP47 antibody (CL488-10694) at dilution of 1:200, CL594-Phalloidin (red).



1X10^6 HeLa cells were intracellularly stained with 0.4 ug CoraLite® Plus 488 Anti-Human TIP47 (CL488-10694) (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).