For Research Use Only

CoraLite® Plus 488-conjugated c-MYC Polyclonal antibody



Purification Method:

IF/ICC: 1:300-1:1200

100 µl suspension

wavelengths:

493 nm / 522 nm

Antigen affinity purification

Excitation/Emission maxima

FC (Intra): 0.40 ug per 10^6 cells in a

Recommended Dilutions:

Catalog Number: CL488-10828

Featured Product

Basic Information

Catalog Number: CL488-10828 BC000141 Concentration: GeneID (NCBI): 1000 μg/ml 4609 **UNIPROT ID:** Source: Rabbit P01106

Full Name: Isotype:

Immunogen Catalog Number:

AG1263

49 kDa Observed MW:

Tested Applications: IF/ICC, FC (Intra)

Species Specificity: human, mouse

GenBank Accession Number:

v-myc myelocytomatosis viral oncogene homolog (avian)

Calculated MW:

62-65 kDa, 50 kDa

Positive Controls:

IF/ICC: HEK-293 cells, FC (Intra): HeLa cells,

Background Information

MYC contains one basic helix-loop-helix (bHLH) domain. This protein is a multifunctional, nuclear phosphoprotein that plays a role in cell cycle progression, apoptosis and cellular transformation. It functions as a transcription factor that regulates transcription of specific target genes. It seems to activate the transcription of growth-related genes. It binds DNA in a non-specific manner, yet also specifically recognizes the core sequence 5'-CAC[GA]TG-3'. Mutations, overexpression, rearrangement and translocation of this gene have been associated with a variety of hematopoietic tumors, leukemias and lymphomas, including Burkitt lymphoma. This antibody is a rabbit polyclonal antibody raised against recombinant protein of human MYC.The 50kDa band recognized by antibody is the native form of MYC, while the other bands, between 60-70kDa, are the phosphorylated form of MYC [PMID: 12189186]

Storage

Applications

Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.


PBS with 50% glycerol, 0.05% Proclin300, 0.5% BSA, pH7.3

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed HEK-293 cells using CoraLite® Plus 488 c-MYC antibody (CL488-10828) at dilution of 1:600.



1X10^6 HeLa cells were intracellularly stained with 0.4 ug CoraLite® Plus 488 Anti-Human c-MYC (CL488-10828) (red), or 0.4 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).