For Research Use Only

CoraLite®488-conjugated FUS/TLS Polyclonal antibody

Catalog Number:CL488-11570 Featured Product



Basic Information

Catalog Number: CL488-11570 Concentration: 1000 μg/ml Source:

UNIPROT ID: Rabbit P35637 Full Name: Isotype: fusion (involved in t(12;16) in

malignant liposarcoma) Immunogen Catalog Number:

AG2150

Calculated MW: 75 kDa

Purification Method:

Antigen affinity purification Recommended Dilutions:

IF/ICC: 1:50-1:500

FC (Intra): 0.20 ug per 10^6 cells in a

100 µl suspension

Excitation/Emission maxima

wavelengths: 491 nm / 516 nm

Applications

Tested Applications: IF/ICC, FC (Intra) Species Specificity: human, mouse, rat

Positive Controls:

IF/ICC: HepG2 cells, HeLa cells

FC (Intra): MCF-7 cells,

Background Information

FUS (also named TLS and POMp75) belongs to the RRM TET family. FUS may play a role in the maintenance of genomic integrity; it binds both single-stranded and double-stranded DNA and promotes ATP-independent $annealing\ of\ complementary\ single-stranded\ DNAs\ and\ D-loop\ formation\ in\ superhelical\ double-stranded\ DNA.\ FUS$ is also an RNA-binding protein, and its links to neurodegenerative disease proffer the intriguing possibility that altered RNA metabolism or RNA processing may underlie or contribute to neuron degeneration[PMID: 22640227]. FUS may be a cause of angiomatoid fibrous histiocytoma (AFH) and is implicated in certain forms of amyotrophic lateral sclerosis (ALS) and frontotemporal dementias (FTDs) such as frontotemporal lobar dementia with ubiquitin inclusions (FTLD-U)[PMID: 22640227]. This antibody is a rabbit polyclonal antibody raised against an internal region of human FUS. This antibody is conjugated with CL488, Ex/Em 488 nm/515 nm.

GenBank Accession Number:

BC026062

GeneID (NCBI):


Storage

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.


PBS with 50% glycerol, 0.05% Proclin300, 0.5% BSA, pH7.3

Aliquoting is unnecessary for -20°C storage

Selected Validation Data




1X10^6 MCF-7 cells were intracellularly stained with 0.2 ug Coralite® 488 Anti-Human FUS/TLS (CL488-11570) (red), or 0.2 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using Coralite® 488 FUS/TLS antibody (CL488-11570) at dilution of 1:200.



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using Coralite®488 FUS/TLS antibody (CL488-11570) at dilution of 1:200.



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using Coralite®488 FUS/TLS antibody (CL488-11570) at dilution of 1:200, Coralite®594 TDP-43 (C-terminal) antibody (CL594-12892, red).