For Research Use Only

CoraLite® Plus 488-conjugated VEGFA Polyclonal antibody



Catalog Number: CL488-19003

Featured Product

Basic Information

Catalog Number: GenBank Accession Number:
CL488-19003 BC065522
Concentration: GeneID (NCBI):
1000 µg/ml 7422
Source: UNIPROT ID:
Rabbit P15692

Isotype: Full Name:

IgG vascular endothelial growth factor A
Immunogen Catalog Number: Calculated MW:

AG13500 412 aa, 46 kDa

Purification Method: Antigen affinity purification Recommended Dilutions:

IF/ICC: 1:300-1:1200 FC (Intra): 0.40 ug per 10^6 cells in a

100 µl suspension

FC: 0.40 ug per 10^6 cells in a 100 µl

suspension

Excitation/Emission maxima

wavelengths: 493 nm / 522 nm

Applications

Tested Applications:

IF/ICC, FC (Intra)

Species Specificity:
human, mouse

Positive Controls:

IF/ICC : HeLa cells,
FC (Intra) : NIH/3T3 cells,

Background Information

VEGFA, also named as VEGF or VPF, belongs to the PDGF/VEGF growth factor family. It is a growth factor active in angiogenesis, vasculogenesis and endothelial cell growth. VEGFA induces endothelial cell proliferation, promotes cell migration, inhibits apoptosis and induces permeabilization of blood vessels. It binds to the FLT1/VEGFR1 and KDR/VEGFR2 receptors, heparan sulfate and heparin. Defects in VEGFA are associated with microvascular complications of diabetes type 1 (MVCD1). VEGFA has 17 isoforms with MW from 16 to 45 kDa. Some isoforms have homodimer forms (e.g.; VEGFA189 38kDa or VEFGA110 34kDa). VEGF-A exists in at least seven homodimeric isoforms. The monomers consist of 121, 145, 148, 165, 183, 189, or 206 amino acids (PMID:15602010). This antibody can recognize all VEGFA isoforms.

Storage


Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer:


PBS with 50% glycerol, 0.05% Proclin300, 0.5% BSA, pH7.3

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunofluorescent analysis of (-20°C Methanol) fixed HeLa cells using CoraLite® Plus 488 VEGFA antibody (CL488-19003) at dilution of 1:600.



1X10^6 NIH/3T3 cells were intracellularly stained with 0.4 ug Coralite® Plus 488 Anti-Human VEGF (CL488-19003) (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).