#### For Research Use Only

# CoraLite® Plus 488-conjugated Ki-67 Polyclonal antibody

proteintech®
Antibodies | ELISA kits | Proteins
www.ptglab.com

Catalog Number: CL488-27309

Featured Product

2 Publications

**Basic Information** 

Catalog Number: CL488-27309 Concentration: 1000 ug/ml Source:

Rabbit Isotype: IgG

Immunogen Catalog Number:

AG26266

GenBank Accession Number:

NM\_002417 GenelD (NCBI): 4288

antigen identified by monoclonal

antibody Ki-67

Calculated MW: 359 kDa

Full Name:

Purification Method: Antigen affinity purification

Recommended Dilutions:

IF/ICC: 1:50-1:500 FC (Intra): 0.80 ug per 10^6 cells in a

100 µl suspension

Excitation/Emission maxima

wavelengths: 493 nm / 522 nm

**Applications** 

Tested Applications: IF/ICC, FC (Intra)
Cited Applications:

IF

Species Specificity:

human Cited Species: human Positive Controls:

IF/ICC: HeLa cells,

FC (Intra): Ramos cells, HeLa cells

### **Background Information**

The Ki-67 protein (also known as MKI67) is a cellular marker for proliferation. Ki67 is present during all active phases of the cell cycle (G1, S, G2 and M), but is absent in resting cells (G0). Cellular content of Ki-67 protein markedly increases during cell progression through S phase of the cell cycle. Therefore, the nuclear expression of Ki67 can be evaluated to assess tumor proliferation by immunohistochemistry. It has been demonstrated to be of prognostic value in breast cancer. In head and neck cancer, several studies have reported an association between high proliferative activity and poorer prognosis.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Ze Dai	38994159	World J Gastrointest Oncol	IF
Yuanhui Xu	38050438	J Biochem Mol Toxicol	IF

Storage

Storage:


Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer


PBS with 50% glycerol, 0.05% Proclin300, 0.5% BSA, pH7.3  $\,$ 

Aliquoting is unnecessary for -20°C storage

## Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using CoraLite® Plus 488 KI67 antibody (CL488-27309) at dilution of 1:200.



1x10^6 Ramos cells were intracellularly stained with 0.8 ug Coralite® Plus 488 Anti-Human Kl67 (CL488-27309)(red), or 0.8 ug Coralite® Plus 488-conjugated Rabbit IgG control Rabbit PolyAb (CL488-30000) (blue). Cells were fixed and permeabilized with True-Nuclear Transcription Factor Buffer Set.