For Research Use Only

## CoraLite® Plus 488-conjugated VDAC1/Porin Polyclonal antibody



Catalog Number:CL488-55259 Featured Product

**Basic Information** 

Catalog Number:
CL488-55259

NM\_003374

Concentration:
GeneID (NCBI):
7416

Source:
UNIPROT ID:
Rabbit
P21796

Isotype:
Full Name:

G voltage-dependent anion channel 1

Calculated MW: 31 kDa Observed MW: 31 kDa Purification Method: Antigen affinity purification Recommended Dilutions: IF-P: 1:50-1:500

FC (Intra): 0.80 ug per 10^6 cells in a 100 µl suspension

Excitation/Emission maxima wavelengths: 493 nm / 522 nm

**Applications** 

Tested Applications: IF-P, FC (Intra) Species Specificity: human, mouse, rat Positive Controls:

IF-P: human liver cancer tissue, FC (Intra): HepG2 cells,

## **Background Information**

VDAC1, also named as VDAC, porin 31HM, porin 31HL and plasmalemmal porin, belongs to the eukaryotic mitochondrial porin family. It adopts an open conformation at low or zero membrane potential and a closed conformation at potentials above 30-40 mV, to form a channel through the mitochondrial outer membrane and also the plasma membrane. Unlike other membrane transport proteins, porins are large enough to allow passive diffusion. Studies have shown that VDAC1 is subject to both phosphorylation and acetylation (PMID: 23233904). The apparent molecular weight of VDAC1 is 30-37 kDa (PMID: 14573604; 23754752; 25681439). Hypoxic conditions were found to trigger cleavage of the VDAC1 C-terminal to yield a 26-kDa truncated but active form (PMID: 22389449; 23233904). This antibody is specific to VDAC1.

Storage


Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer:


PBS with 50% glycerol, 0.05% Proclin300, 0.5% BSA, pH7.3

Aliquoting is unnecessary for -20°C storage

## Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed human liver cancer tissue using Coralite® Plus 488 VDAC1/Porin antibody (CL488-55259) at dilution of 1:200.



1x10^6 HepG2 cells were intracellularly stained with 0.8 ug Coralite® Plus 488-conjugated VDAC1/Porin Polyclonal antibody (CL488-55259) (red), or 0.8 ug Coralite® Plus 488-conjugated Rabbit IgG control Rabbit PolyAb (CL488-3000) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).