For Research Use Only

CoraLite® Plus 488 Anti-Mouse CD16/CD32 (2.4G2)

Catalog Number: CL488-65080



Basic Information

Catalog Number:GenBank Accession Number:Purification Method:CL488-65080BC038070Affinity purification

 Concentration:
 GeneID (NCBI):
 CloneNo.:

 100ug, 0.5 mg/ml
 14130
 2.4G2

Source: UNIPROT ID: Recommended Dilutions: Rat P08101 FC: 0.25 ug per 10^6 cells in 100 $\,\mu$ l

Isotype: Full Name: suspension

IgG2b, kappa Fc receptor, IgG, low affinity IIb Excitation/Emission maxima

wavelengths: 493 nm / 522 nm

Applications

Tested Applications:

FC

Species Specificity:

mouse

Positive Controls:

FC: mouse splenocytes,

Background Information

CD16 (Fc γ RIII) is a 50-70 kDa low affinity Fc receptor found on the surface of natural killer cells, neutrophil polymorphonuclear leukocytes, monocytes and macrophages. CD16 mediates antibody-dependent cellular cytotoxicity (ADCC) and other antibody-dependent responses, such as phagocytosis. CD32 (Fc γ RII) is a 40 kD transmembrane glycoprotein that binds to the Fc region of IgG with low affinity. CD32 is present on phagocytic cells such as macrophages and neutrophils, and is involved in the process of phagocytosis and clearing of immune complexes. The 2.4G2 antibody reacts with the extracellular domains of mouse CD16 (FcRIII) and CD32 (FcRII) (PMID: 90108; 8406898).

Storage


Storage:

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

torage Buffer

PBS with 0.09% sodium azide and 0.5% BSA, pH7.3

Selected Validation Data



1X10^6 mouse splenocytes were surface stained with 0.25 ug CoraLite® Plus 488 Anti-Mouse CD16 / CD32 (CL488-65080, Clone:2.4G2) (red) or 0.25 ug Control Antibody. Cells were not fixed.