For Research Use Only

## CoraLite® Plus 488-conjugated Cytokeratin 18 Monoclonal antibody



Catalog Number: CL488-66187

**Basic Information** 

Catalog Number: GenBank Accession Number: CL488-66187 BC000180

CL488-66187 BC000180 Protein G purification
Concentration: Genel D (NCBI): CloneNo.:

1000  $\mu$  g/ml 3875 1G11C4 Source: UNIPROT ID: Recommended Dilutions:

Mouse P05783 IF/ICC: 1:50-1:500

Isotype:Full Name:FC (Intra): 0.50 ug per 10^6 cells in aIgG1keratin 18100 μl suspension

FC: 0.50 ug per 10^6 cells in a 100 µl
Immunogen Catalog Number: Calculated MW:

AG1260 48 kDa suspension

Excitation/Emission maxima wavelengths:

wavelengths: 493 nm / 522 nm

**Purification Method:** 

**Applications** 

Tested Applications: IF/ICC, FC (Intra)

Species Specificity: human, canine

Positive Controls:

IF/ICC: HeLa cells, HepG2 cells, MDCK cells

FC (Intra): HeLa cells,
FC: HeLa cells.

## **Background Information**

Keratins are a large family of proteins that form the intermediate filament cytoskeleton of epithelial cells, which are classified into two major sequence types. Type I keratins are a group of acidic intermediate filament proteins, including K9-K23, and the hair keratins Ha1-Ha8. Type II keratins are the basic or neutral courterparts to the acidic type I keratins, including K1-K8, and the hair keratins, Hb1-Hb6. KRT18, also named as CYK18, PIG46 and K18, is the most commonly found members of the intermediate filament gene family. KRT18 is a type I keratin which is found primarily in non squamous epithelia and is present in a majority of adenocarcinomas and ductal carcinomas but not in squamous cell carcinomas. The antibody is conjugated with CL488, Ex/Em 488 nm/515 nm.

Storage

Storage:


Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer


PBS with 50% glycerol, 0.05% Proclin300, 0.5% BSA, pH7.3

Aliquoting is unnecessary for -20°C storage

## Selected Validation Data



Immunofluorescent analysis of (-20°C Ethanol) fixed HeLa cells using CL488-66187 (Cytokeratin 18 antibody) at dilution of 1:100.



1X10^6 HeLa cells were intracellularly stained with 0.5 ug CoraLite® Plus 488 Anti-Human Cytokeratin 18 (CL488-66187, Clone:1G11C4) (red), or 0.5 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).