For Research Use Only

# CoraLite®555-conjugated Caspase 9/P35 Polyclonal antibody

www.ptglab.com

**Purification Method:** 

100 ul suspension

wavelengths:

Antigen affinity purification

Excitation/Emission maxima

FC (Intra): 0.40 ug per 10^6 cells in a

Recommended Dilutions:

Catalog Number:CL555-10380

Featured Product

**Basic Information** 

Catalog Number: CL555-10380 Concentration:

1000 ug/ml Source:

Rabbit Isotype:

Immunogen Catalog Number:

AG0404

**Tested Applications:** FC (Intra) Species Specificity:

human, mouse, rat

GenBank Accession Number:

BC002452 GeneID (NCBI):

**UNIPROT ID:** P55211

peptidase

Full Name: caspase 9, apoptosis-related cysteine 557 nm / 570 nm

Calculated MW: 46 kDa

Observed MW: 46 kDa, 35 kDa

Positive Controls:

FC (Intra): HepG2 cells,

## **Background Information**

**Applications** 

Caspase 9, apoptosis-related cysteine protease (CASP9, synonyms: MCH6, APAF3, APAF-3, ICE-LAP6, CASPASE-9c)is a member of the cysteine-aspartic acid protease (caspase) family. Sequential activation of caspases plays a central role in the execution-phase of cell apoptosis. Caspases exist as inactive proenzymes which undergo proteolytic processing at conserved aspartic residues to produce 2 subunits, large and small, that dimerize to form the active enzyme. Caspase 9 is processed by APAF1; this step is thought to be one of the earliest in the caspase activation cascade. 10380-1-AP can recognize the pre- and cleaved- caspase 9. In recent years, the localization of caspase9 was a focus of interest. Beside its cytoplasmic distribution, a very extensive localization study was done on rat brain tissue, where caspase9 was found located predominantly in the nucleus and to a lesser extend in the cytoplasm [PMID: 15541731].

### Storage


#### Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 50% glycerol, 0.05% Proclin300, 0.5% BSA, pH7.3

Aliquoting is unnecessary for -20°C storage

# Selected Validation Data



1X10^6 HepG2 cells were intracellularly stained with 0.4 ug Coralite®555 Anti-Human Caspase 9/P35 (CL555-10380) (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).