For Research Use Only

CoraLite®555-conjugated c-MYC Polyclonal antibody

Catalog Number: CL555-10828 Featured Product



Basic Information

Catalog Number: GenBank Accession Number: CL555-10828 BC000141 Concentration: GeneID (NCBI): 1000 μg/ml 4609 **UNIPROT ID:** Source: Rabbit P01106

Full Name: Isotype: v-myc myelocytomatosis viral

Immunogen Catalog Number:

AG1263

oncogene homolog (avian) Calculated MW: 49 kDa

> Observed MW: 62-65 kDa, 50 kDa

Purification Method:

Antigen affinity purification Recommended Dilutions:

FC (Intra): 0.20 ug per 10^6 cells in a

100 µl suspension

IF/ICC: 1:50-1:500

Excitation/Emission maxima wavelengths:

557 nm / 570 nm

Applications

Tested Applications: Positive Controls: IF/ICC, FC (Intra) IF/ICC: HEK-293 cells, Species Specificity: FC (Intra): Raji cells, human, mouse

Background Information

MYC contains one basic helix-loop-helix (bHLH) domain. This protein is a multifunctional, nuclear phosphoprotein that plays a role in cell cycle progression, apoptosis and cellular transformation. It functions as a transcription factor that regulates transcription of specific target genes. It seems to activate the transcription of growth-related genes. It binds DNA in a non-specific manner, yet also specifically recognizes the core sequence 5'-CAC[GA]TG-3'. Mutations, overexpression, rearrangement and translocation of this gene have been associated with a variety of hematopoietic tumors, leukemias and lymphomas, including Burkitt lymphoma. This antibody is a rabbit polyclonal antibody raised against recombinant protein of human MYC.The 50kDa band recognized by antibody is the native form of MYC, while the other bands, between 60-70kDa, are the phosphorylated form of MYC [PMID: 12189186]

Storage

Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.


PBS with 50% glycerol, 0.05% Proclin300, 0.5% BSA, pH7.3

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed HEK-293 cells using CoraLite®555 c-MYC antibody (CL555-10828) at dilution of 1:200.



1X10^6 Raji cells were intracellularly stained with 0.2 ug CoraLite® 555 Anti-Human c-MYC (CL555-10828) (red), or 0.2 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).