For Research Use Only

CoraLite®555-conjugated MECP2 Polyclonal antibody

Catalog Number:CL555-10861 Featured Product



Basic Information

Catalog Number: GenBank Accession Number:
CL555-10861 BC011612
Concentration: GeneID (NCBI):
1000 µg/ml 4204
Source: UNIPROT ID:
Rabbit P51608

Isotype: Full Name:
IgG methyl CpG binding protein 2 (Rett

Immunogen Catalog Number: syndrome)
AG1309 Calculated MW:

52 kDa, 53 kDa Observed MW: 75 kDa Purification Method: Antigen Affinity Purified Recommended Dilutions:

IF/ICC: 1:50-1:500 FC (Intra): 0.20 ug per 10^6 cells in a

100 µl suspension

Excitation/Emission maxima

wavelengths: 557 nm / 570 nm

Applications

Tested Applications:

IF/ICC, FC (Intra)

Species Specificity:
human, mouse

Positive Controls:

IF/ICC : HepG2 cells,

FC (Intra) : NIH/3T3 cells,

Background Information

METHYL-CpG-BINDING PROTEIN 2(MECP2), is a chromatin-associated protein that can both activate and repress transcription. Mecp2 in the adult mouse is high in the brain, lung, and spleen, lower in heart and kidney. MECP2 takes part in the control of neuronal activity-dependent gene regulation, and this process may underlie the pathology of Rett syndrome, a severe developmental disorder with autistic phenotypes. As compared to wild-type monkeys, MECP2 transgenic monkeys exhibited a higher frequency of repetitive circular locomotion and increased stress responses, as measured by the threat-related anxiety and defensive test. Chromosomal protein that binds to methylated DNA. It can bind specifically to a single methyl-CpG pair. It is not influenced by sequences flanking the methyl-CpGs. Mediates transcriptional repression through interaction with histone deacetylase and the corepressor SIN3A. Two isoforms of MECP2 exist due to alternative splicing events. This antibody reacts with the MECP2 and phosphorylated MECP2 proteins. The calculated molecular weight of MECP2 is a 52 kDa, but the post-modified MECP2 protein is about 75-80 kDa (PMID: 12160743).

Storage

Storage:


Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer


PBS with 50% glycerol, 0.05% Proclin300, 0.5% BSA, pH7.3

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



1X10^6 NIH/3T3 cells were intracellularly stained with 0.2 ug Coralite®555 Anti-Human MECP2 (CL555-10861) (red), or 0.2 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using CoraLite®555 MECP2 antibody (CL555-10861) at dilution of 1:100.