For Research Use Only

CoraLite®555-conjugated Galectin-3 Monoclonal antibody



Catalog Number: CL555-60207

Featured Product

Basic Information

Catalog Number: GenBank Accession Number: Purification Method: CL555-60207 BC001120 Protein A purification

 Source:
 UNIPROT ID:
 Recommended Dilutions:

 Mouse
 P17931
 IF/ICC: 1:50-1:500

Isotype: Full Name: FC (Intra): 0.80 ug per 10^6 cells in a

IgG2b lectin, galactoside-binding, soluble, 3 100 µl suspension

Immunogen Catalog Number: Calculated MW: Excitation/Emission maxima

AG6753 26 kDa wavelengths: 557 nm / 570 nm

Observed MW: 31 kDa

Applications

Tested Applications:

IF/ICC, FC (Intra)

Species Specificity:
human, mouse, rat

Positive Controls:

IF/ICC : A431 cells,

FC (Intra) : HeLa cells,

Background Information

Galectins are a family of animal lectins defined by shared characteristic amino-acid sequences and affinity for β -galactose-containing oligosac-charides (PMID: 8063692). Galectin-3, a member of the β -galactoside-binding proteins, contains one carbohydrate recognition domain (CRD) and a proline- and glycine-rich N-terminal domain through which is able to form oligomers (PMID: 14758078). Galectin-3 is widely expressed in many normal tissues and a variety of tumors. It is found intracellularly in nucleus and cytoplasm or secreted outside of cell, being present on the cell surface or in the extracellular space (PMID: 16478649). Galectin-3 is involved in various biological processes including cell growth, adhesion, differentiation, apoptosis, angiogenesis, immune response, neoplastic transformation and metastasis (PMID: 16478649; 14758078).

Storage


Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer:


PBS with 50% glycerol, 0.05% Proclin300, 0.5% BSA, pH7.3

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunofluorescent analysis of (-20°C Methanol) fixed A431 cells using CoraLite® 555 Galectin-3 antibody (CL555-60207, Clone: 1C1B2) at dilution of 1:200.



1X10^6 HeLa cells were intracellularly stained with 0.8 ug CoraLite®555 Anti-Human Galectin-3 (CL555-60207, Clone:1C 1B2) (red), or 0.8 ug Isotype Control. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).